



# TIER 2 DAY 2

## OCTOBER 28, 2015

Center for Deployment Psychology  
Uniformed Services University of the Health Sciences





## Ethical Considerations for Working with Military Members and Veterans

Center for Deployment Psychology  
Uniformed Services University of the Health Sciences



## Disclaimer

The views expressed are those of the presenters and do not necessarily reflect the opinions of the Uniformed Services University of the Health Sciences, the Department of Defense, or the U.S. Government.



## Learning Objectives

1. Define ethics as it relates to the role of the mental health provider.
2. Identify five (5) ethical challenges common to mental health providers working with the military population.
3. Discuss Gottlieb's model for avoiding dual relationships.
4. Demonstrate knowledge of the ethical decision making process through interactive discussion of military case examples during the presentation.



## What is Ethical



The Martian Agency (2010)



5



## Why Ethics?

We do not act rightly because we have virtue or excellence, but we rather have those because we have acted rightly.

- Aristotle, 384-322 B.C., Greek philosopher and scientist, student to Plato

Even the most rational approach to ethics is defenseless if there isn't the will to do what is right.

- Alexander Solzhenitsyn, Author, winner of the 1970 Nobel Prize for Literature

**You WILL be exposed to ethical dilemmas**



6



## Questions about Ethics



Values are like fingerprints. Nobody's are the same, but you leave 'em all over everything you do.  
— Elvis Presley



7

(DoD photo by Mass Communication Specialist 3rd Class Jasmine Sheard, U.S. Navy/Released)



## What are Ethics?



8

U.S. Navy photo by Chief Mass Communication Specialist Tiffini Jones Vanderwyst/Released



## Ethics?

Merriam Webster's Dictionary offers the following definitions

*Ethics:*

1. A treatise on morals. 2. The science of moral duty; broadly, the science of ideal human character. 3. Moral principles, quality, or practice.

*Ethical:*

1. Of or relating to moral action, motive, or character; also, treating of morals, or ethics. 2. Conforming to professional standards of conduct.

*But as an action...*

- Thinking about reasons in terms of values in a manner that is open to public scrutiny



9



## Ethics

- Therapists must integrate their personal ethical and value traditions with psychology's

- Two major variables:

- 1) Maintenance refers to the degree that we retain the ethical and value traditions of our culture of origin

- 2) Contact and participation refers to the degree to which new psychologists adopt the traditions, norms, values of their new professional culture

Handelsman et al (2005) in Gottlieb et al (2008)



10



## Ethics Standards Are Not Enough

Principle ethics: Obligations to consider when deciding "what to do."

Virtue ethics: Ideals to consider when deciding "who shall I be?"

Principle A: "Psychologists strive to benefit those they serve and take care to avoid harm."

Never let your sense of morals prevent you from doing what's right.  
— Isaac Asimov

Meara et al (1996)



11



## Is It Ethical?



Campbell et al. (Writers), & Leddy (Director) (2001)



12



## Ethics Standards Are Not Enough

A Psychologist with virtues and principles is one who is:

- Motivated to do good
- Possesses vision and discernment
- Emotionally intelligent
- Self-aware
- Appreciates and respects community mores in decision-making

You can easily judge the character of a man by how he treats those who can do nothing for him.  
— James D. Miles



13



## Ethics Continued: Profession Specific

- General Principles: are considered aspirational in nature and are intended to be considered when confronting ethical dilemma's
- Ethical Standards: are purposely written broadly to apply to psychologists in varied roles and the particular application of a standard can vary depending on the context
- Ethics must be practical:
  - "Every clinician is unique – every client is unique."
  - "Ethics that are out of touch with the practicalities of clinical work... are useless."

Pope & Vasquez (1998) in Barnett et al (2007)



14



## Dilemma vs. Conflict



15

U.S. Navy photo by Mass Communication Specialist 2nd Class Michael Russell/Released



## What is it?

### Ethical Dilemma

An ethical dilemma is a situation in which conforming to professional standards of conduct creates a need to make a choice between equally unsatisfactory alternatives. This often results from a discrepancy between professional ethics and law or institutional policy.

### Ethical Conflict

For an ethical–legal discrepancy to become a conflict, the provider's obligations under the law and the provider's obligations under his or her professional code of ethics must be mutually exclusive.



16

Johnson et al (2010), pg. 549



## Conflict: DoD Policy and APA Ethics Code


Psychologists in a military setting face challenges with informed consent:

- Military mission
- Who is the client?
- The impact on the SM's career



## Strategic Model for Managing Ethical Concerns

"Hope is not a plan"



## Forensic Psychology Ethical Decision Making Model

1. Identify the problem
2. Consider the significance of the context and setting
3. Identify and utilize ethical and legal resources
4. Consider personal beliefs and values
5. Develop possible solutions to the problem
6. Consider the potential consequences of various solutions
7. Choose and implement a course of action
8. Assess the outcome and implement changes as needed



## Considerations for Discussion

- I. Boundaries of Competence
- II. Informed Consent
- III. Disposition Driven Diagnosis
- IV. Multiple Relationships
- V. Professional's Own Fitness



## I. Boundaries of Competence

- This is a unique population with its own cultural identity. Is the therapist aware of this culture?
- Is the therapist trained to treat problems and disorders common to military members and veterans?
- Standard 2.01 (Boundaries of Comp.)
- Standard 2.02 (Emergencies)



## Boundaries of Competence

- Military Culture
  - Language
  - Demographics
  - Rank and organizational structure
  - Manners and normative behaviors
  - Beliefs, mission, and values



## Increase Military Cultural Competency

- |                                      |                                      |
|--------------------------------------|--------------------------------------|
| I. Make a Self Inventory             | I. Exposure to Military Culture      |
| II. Adapt Care to Military Culture   | II. Training on Military Regulations |
| III. Attend Military Activities      | III. Training Through Observation    |
| IV. Increase Off-Post Social Support |                                      |



## Boundaries of Competence: The Marine Case #1




## II. Informed Consent



## Informed Consent: The National Guard Student Case #2



## III. Disposition Drives Diagnosis



## Problems with Administratively Driven Diagnoses

“Psychologists base the opinions contained in their recommendations, reports, and diagnostic evaluative statements on information and techniques sufficient to substantiate their findings.”

APA (2010)







## Problems with Administratively Driven Diagnoses

- The “psychologist as administrative broker” role can have unintended consequences for service members.
  - Can perpetuate view of psychologists and mental health diagnosis as imprecise and psychiatric disorders as meaningless or silly.
  - Can also can lead to increased stigma for seeking treatment and devaluing of psychological services



## Problems with Administratively Driven Diagnoses

We are ethically obligated to provide correct diagnosis no matter how the chips fall...

- “My wife says she will leave me/I will lose my job if I have to deploy again.”
- “I have to go home right now to save my marriage.”
- “Is it really MDD?”



## Hot-Button Issues

- Can create lots of tension between the patient and the military with conflicting goals for each side
- Can be popular diagnoses that get lots of attention in the media and may be misunderstood or stereotyped in popular culture



## Hot-Button Issues

- Substance abuse
- PTSD
- Suicidal behavior
- Who is the client
- Implications of diagnoses
- How do you balance the following:
  - Potential secondary gain
  - Stigma
  - Barriers to care
  - Confidentiality



## Disposition Drives Diagnosis: “Off the Record” Case #3



## IV. Multiple Relationships

- Military members can often present opportunities to create dual relationship.
- You must approach them carefully and thoughtfully.



## Multiple Relationships

**Multiple Relationships occur when a psychologist:**

- Has more than one role with a client
- Has a relationship with a person closely associated with the client
- Is likely to enter into another relationship with the client in the future



## Gottlieb’s Five-Step Model for Avoiding Dual Relationships

**Step 1.** Assess the current relationship according to three dimensions:

1. Power of provider in the relationship
2. Duration of the existing relationship
3. Termination is clearly defined and definitive

**Step 2.** Look at the contemplated (dual) relationship from the three dimensions.

If these three dimensions are all high, the relationship should be avoided because there is risk for harm (high therapist power, long term relationship and no clear/specified termination).



## Gottlieb's Five-Step Model for Avoiding Dual Relationships

**Step 3.** When the three dimensions fall in the mid-range, examine both relationships for incompatible roles.

**Step 4.** Obtain consultation from a colleague.

**Step 5.** Discuss the decision with the consumer/patient.

Dual relationships may be even more complicated for military providers working at a military treatment facility

Gottlieb (1993)



40



## Multiple Relationships: Petty Officer 3<sup>rd</sup> Class Case #4



41



## V. Professional's Own Fitness



43



## Professional's Own Fitness Personal Problems & Conflicts

### Standard 2.06

- Psychologists refrain from activities when they know or should know that there is a substantial likelihood that their personal problems will prevent them from performing their work-related activities in a competent manner.
- When impaired, they take appropriate measures to obtain professional help and determine whether they should limit or suspend practice.

American Psychological Association (2010)



44



## Professional's Own Fitness Personal Problems & Conflicts

- Exposed to stories of human suffering, war and death
- Accountable to engage in self-assessment
- Accountable to seek assistance
- Accountable to scrutinize the fitness of colleagues
- Any difficulties here?



## Professional's Own Fitness: Skipping Lunch and Scrambling Case #5



## Ethics in Action



## Ethics in Action





## A Reminder

- Standard 1.02: Conflicts between Ethics and Law, Regulations, or other Governing Authority.

If psychologist's ethical responsibilities conflict with law, regulations, etc., psychologists make known their commitment to the Ethics Code...If the conflict is irresolvable, psychologists **may** adhere to the requirements of the law...

*Remember to be vigilant of self and other providers to ensure ethical and safe practice.*



## Key Points

Examine several models for anticipating and responding to ethical dilemmas:

- Utilize a structured decision making model
- Talk to peers/colleagues
- Have a list of experts for consultation
- Document discussions & actions
- Be mindful of behavioral drift
- Decision-making in ethics always involves a process
  - This process involves thinking about values
- Good law, good ethics, and good clinical care go hand-in-hand
- Never worry alone



## Summary

- Personal/professional ethics
- The decision-making process
- Strategies for anticipating and responding to ethical dilemmas
- Ethical dilemmas encountered by clinicians working with military members and veterans
- How the APA Ethics Code may conflict with DoD regulations, law and/or policy



## CDP Website: Deploymentpsych.org

### Features include:

- Descriptions and schedules of upcoming training events
- Blog updated daily with a range of relevant content
- Articles by subject matter experts related to deployment psychology, including PTSD, mTBI, depression, and insomnia
- Other resources and information for behavioral health providers
- Links to CDP's Facebook page and Twitter feed





## Online Learning

The following online courses are located on the CDP's website at:

[Deploymentpsych.org/training/online-courses](http://Deploymentpsych.org/training/online-courses)

**NOTE: All of these courses can be taken for free or for CE Credits for a fee**

- Cognitive Processing Therapy (CPT) for PTSD in Veterans and Military Personnel (1.25 CE Credits)
- Prolonged Exposure Therapy for PTSD in Veterans and Military Personnel (1.25 CE Credits)
- Epidemiology of PTSD in Veterans: Working with Service Members and Veterans with PTSD (1.5 CE Credits)
- Provider Resiliency and Self-Care: An Ethical Issue (1 CE Credit)
- Military Cultural Competence (1.25 CE Credits)
- The Impact of Deployment and Combat Stress on Families and Children, Part 1 (2.25 CE Credits)
- The Impact of Deployment and Combat Stress on Families and Children, Part 2 (1.75 CE Credits)
- The Fundamentals of Traumatic Brain Injury (TBI) (1.5 CE Credits)
- Identification, Prevention, & Treatment of Suicidal Behavior in Service Members & Veterans (2.25 CE Credits)
- Depression in Service Members and Veterans (1.25 CE Credits)

*All of these courses and several others are contained in the Serving Our Veterans Behavioral Health Certificate program, which also includes 20+ hours of Continuing Education Credits for \$350.*



54



## Provider Support

CDP's "Provider Portal" is exclusively for individuals trained by the CDP in evidence-based psychotherapies (e.g., CPT, PE, and CBT-I)

Features cover topics including:

- Consultation message boards
- Hosted consultation calls
- Printable fact-sheets, manuals, handouts, and other materials
- FAQs and one-on-one interaction with answers from SMEs
- Videos, webinars, and other multimedia training aids

Participants in CDP's evidence-based training will automatically receive an email instructing them how to activate their user name and access the "Provider Portal" section at [Deploymentpsych.org](http://Deploymentpsych.org).



55



## How to Contact Us

### Center for Deployment Psychology

Department of Medical & Clinical Psychology

Uniformed Services University of the Health Sciences

4301 Jones Bridge Road, Executive Office: Bldg. 11300-602

Bethesda, MD 20813-4768

**Email:** [General@DeploymentPsych.org](mailto:General@DeploymentPsych.org)

**Website:** [DeploymentPsych.org](http://DeploymentPsych.org)

**Facebook:** <http://www.facebook.com/DeploymentPsych>

**Twitter:** @DeploymentPsych



56



References

- Alda, A., Metcalfe, B., Rappaport, J., Wilcox, D., Mumford, T., Davis, E., Pollock, D., & Hall, K. (Writers), & Alda, A. (Director). (1983). *M.A.S.H.: Goodbye, Farewell and Amen* [Television Series Episode]. In Metcalfe, B., Rappaport, J., Mumford, T., Wilcox, D., Tischler, S. (Producers), *M.A.S.H.*. Los Angeles, CA: 20<sup>th</sup> Century Fox Television.
- American Psychological Association. (2010). Ethical principles of psychologists and code of conduct. Retrieved from <http://apa.org/ethics/code/index.aspx>
- Barnett, J. E., Behnke, S. E., Rosenthal, S. L., & Koocher, G. P. (2007). In case of ethical dilemma, break glass: Commentary on ethical decision making in practice. *Professional Psychology: Research and Practice*, 38(1), 7-12. <http://dx.doi.org/10.1037/0003-066X.64.8.793>
- Bush, S. S., Connell, M. A., & Denney, R. L. (2006). *Ethical practice in forensic psychology; A systematic model for decision making*. Washington, DC: APA Books.
- Campbell, G. Dombrowski, Adams, B., Ascencio, N., et al. (Writers), & Leddy, B. (Director). (2001). *MADtv: Season 6, Episode 24* [Television Series Episode]. In Adams, B., Blasucci, D., Dombrowski, L., Hartt, B., Jones, Q., Salzman, D., & Sites, S. (Producers), *MADtv*. Los Angeles, CA: Fox Television Studios.
- Gottlieb, M. C. (1993). Avoiding exploitive dual relationships: A decision-making model. *Psychotherapy*, 30(1), 41-48. <http://dx.doi.org/10.1037/0033-3204.30.1.41>
- Gottlieb, M. C., Handelsman, M. M., & Knapp, S. (2008). Some principles for ethics education: Implementing the acculturation model. *Training and Education in Professional Psychology*, 2(3), 123-128. <http://dx.doi.org/10.1037/1931-3918.2.3.123>
- Jeffrey, T.B., Rankin, R.J., & Jeffrey, L.K. (1992). In the service of two masters: The ethical-legal dilemma faced by military psychologists. *Professional Psychology: Research and Practice*, 23(9), 91-95.
- Johnson, W. B. (2008). Top ethical challenges for military clinical psychologists. *Military Psychology*, 20(1), 49-62. <http://dx.doi.org/10.1080/08995600701753185>
- Johnson, W. B., Grasso, I., & Maslowski, K. (2010). Conflicts between ethics and law for military mental health providers. *Military Medicine*, 175(8), 548-553.
- Kennedy, C. H., & Johnson, B. W. (2009). Mixed agency in military psychology: Applying the American Psychological Association Ethics Code. *Psychological Services*, 6(1), 22-31. <http://dx.doi.org/10.1037/a0014602>
- Luby, C. D. (2012). Promoting military cultural awareness in off-post community of behavioral health and social support service providers. *Advances in Social Work*, 13(1), 67-82.
- McCauley, M., Hughes, J.H., & Liebling-Kalifani, H. (2008). Ethical Considerations for military clinical psychologists: A review of selected literature. *Military Psychology*, 20, 7-20. <http://dx.doi.org/10.1080/08995600701753128>



- Meara, N. M., Schmidt, L. D., & Day, J. D. (1996). Principles and virtues: A foundation for ethical decisions, policies, and character. *The Counseling Psychologist, 24*(1), 4-77. <http://dx.doi.org/10.1177/0011000096241002>
- Orme, D. R., & Doerman, A. L. (2001). Ethical dilemmas and U.S. Air Force clinical psychologists: A Survey. *Professional Psychology: Research and Practice, 32*(3), 305-311. <http://dx.doi.org/10.1037//0735-7028.32.3.305>
- Reger, M. A., Etherage, J. R., Reger, G. M., & Gahm, G. A. (2008). Civilian psychologists in an Army culture: The ethical challenge of cultural competence. *Military Psychology, 20*(1), 21-35. <http://dx.doi.org/10.1080/08995600701753144>
- Staal, M. A., & King, R. E. (2000). Managing a multiple relationship environment: The ethics of military psychology. *Professional Psychology: Research and Practice, 31*(6), 698-705. <http://dx.doi.org/10.1037//0735-7028.31.6.698>
- The Martian Agency (Producers). (2010). Does a former drill sergeant make a terrible therapist? [Television commercial]. Richmond, VA: GEICO Insurance Company.

Boundaries of Competence:

The Marine

Case #1

You are one of two civilian psychologists working in a local clinic in a small town with no nearby military treatment facilities or VA clinics. Neither you nor the other provider have prior experience treating service members. A client is referred to you who is a 28-year-old, divorced, Caucasian male. The intake notes from the referral indicate that he was a Marine for 9 years with three combat deployments: two for OIF in 2007 and one for OEF in 2009. According to the intake notes, his divorce was finalized last year; he had his second DUI last month; and during deployment, several of his buddies were killed in combat when their vehicle blew up, which he witnessed, and another buddy completed suicide during the past year. Based on the intake notes the client denied any suicidal ideation but reported complaints that could suggest PTSD like nightmares, bouts of irritability, disturbed sleep, avoidance of reminders of his buddies' deaths, seeing images of the burning vehicle, and feeling “numb” and disconnected from others.

---

Informed Consent:

The National Guard Student

Case #2

You are a civilian social worker in a university counseling center treating a 20-year-old, Caucasian female in the National Guard. She entered counseling because of disciplinary measures stemming from an underage drinking ticket, her first legal complication. In your third session, you learn that she is scheduled to deploy in support of Operation Enduring Freedom (OEF – Afghanistan) after the semester. In the intake, she had indicated that she drinks regularly with her friends (mostly weekends), and that she has blacked out several times at football tailgate parties. During this session, when you ask her more about her National Guard experience and what her MOS (job) is, she tells you she is qualified to drive trucks and will likely be driving outside the wire delivering supplies as part of a convoy. She states that she has good friends in the National Guard, and that she drinks with them when she can.

Disposition Drives Diagnosis:  
“Off the Record”  
Case #3

You are a psychologist in private practice and accept Tri-Care. You have an appointment with a 23-year-old, single, African American male who is a four year Army veteran. He states he recently separated from the Army in order to go to school, but that he intends to re-enter the military and attend Officer’s Candidate School (OCS) when he completes his degree. He is hoping to have a 20-year military career and to be in Special Forces. He denies any history of psychiatric treatment, but indicates that he is seeking your assistance because he has periods of time when he rarely sleeps and yet can be incredibly productive during these times; recently he has been able to read ahead of his classmates, and when in the Army he “never slept much anyway.” However, he has been getting into fights recently. Also, his girlfriend broke up with him after he drove to Alabama (a 2-day drive) to see an Army buddy that he wanted to have a beer with. He’s speaking very quickly and his stories can be tangential, but he can be redirected when asked direct questions. During the interview, he interrupts himself to say, “By the way, I don’t want you to write anything in the notes. I just have to talk this stuff out. I came to you so that this wouldn’t go into my record. That’s why I’m going to pay you in cash.” He explains that he won’t be using his Tri-Care benefits.

---

Multiple Relationships:  
Petty Officer 3<sup>rd</sup> Class  
Case #4

You are a Military OneSource Provider in a city near a large Naval base. One of your patients is a 34-year-old, Asian American Lieutenant Commander JAG (Naval lawyer), whom you are seeing for mild depression and anxiety related to work stress. Another of your patients is a 22-year-old Caucasian female Petty Officer 3rd Class (PO3) who is experiencing distress regarding a history of childhood abuse and a recent military sexual assault, which she has not reported to legal services. One day, the JAG calls you and tells you that he is calling as the legal representative for his command, which is also the command of the PO3. He says that the PO3 was hospitalized last weekend after a breakdown on her ship in which she threatened to kill herself or someone else. He says the hospital psychiatrist recommended she be administratively separated for a personality disorder. The JAG tells you that he is handling this case and that the PO3 is fighting the separation. She has requested a formal hearing (administrative separation board) and asked him to contact you to confirm her earlier report of sexual assault. She has also requested you testify at the separation board.

---

Professional's Own Fitness:  
Skipping Lunch and Scrambling  
Case #5

You are a therapist working for at a local women's health clinic. You have become reputable as one of few providers in the area and one who successfully treats women with symptoms related to military sexual trauma. You find yourself skipping lunch, staying at work late to finish notes, avoiding exercise because you are so exhausted, and grimacing frequently with back pain. Every once in a while, you have trouble focusing on your clients during sessions. Twice a week you scramble to make it to an evening class on treating eating disorders in women- an area you are less familiar with but encounter in the clinic - because this seminar is rarely offered where you live. Today you had to leave work early when your 15-year-old niece who is staying with you for the summer called because she had gotten in a car accident.





**CDP CENTER FOR  
DEPLOYMENT PSYCHOLOGY**  
*Preparing Professionals to Support Warriors and Families*



## **Assessment and Treatment of Depression and Suicidal Behavior Associated with Military Service: An Overview**

Center for Deployment Psychology  
Uniformed Services University of the Health Sciences



## **Disclaimer**

The views expressed are those of the presenters and do not necessarily reflect the opinions of the Uniformed Services University of the Health Sciences, the Department of Defense, or the U.S. Government.



## **Learning Objectives**

1. Discuss the prevalence of depression and suicide in the military population.
2. Describe the correct nomenclature for suicidal and related behaviors.
3. Identify strategies for screening and assessing military clients for depression and suicidal behaviors.
4. Review effective therapies for treating military clients with depression and those displaying suicidal behaviors.



- What are your negative thoughts about seeing suicidal clients in general?
- What are/could be the hardest parts about working with suicidal military or veteran clients?



## Outline

- Military depression and suicide rates
- Etiology of depression and suicide
- Depressive Spectrum Disorders: diagnostic criteria
- Suicide risk factors, warning signs & protective factors
- Assessment of depression and suicide
- Treatment of depression and suicidal behavior



## Military Health Significance of Depression and Suicide



## Depression in Returning OIF/OEF Service Members

12 Month Post-Deployment	Depression Symptoms	Depression Symptoms/ Some Impairment	Depression Symptoms/ Functional Impairment
Active Component	15.7%	14.4%	8.5%
Reserve Component	15.9%	13.7%	7.3%



## Depression and Deployment in Millennium Cohort Study

### New Onset Depression

	Men	Odds Ratio	Women	Odds Ratio
Never Deployed	3.9%	1.0	7.7%	1.0
Deployed, No Combat	2.3%	.66	5.1%	.65
Deployed, Combat	5.7%	1.32	15.7%	2.13

Wells et al. (2010)




10



## 2011 DoD Survey: Level of Depression

### 2011 Survey of Health Related Behaviors Among Active Duty Military Personnel



DoD 2011 Health Related Behaviors Survey (2013)



11



## Depression in Veterans

- 14% of veterans are diagnosed with depression  
–Yet it is likely under-diagnosed
- 11% of veterans aged 65+ y/o are diagnosed with MDD (twice the rate of adults 65+ in the general population)

National Alliance on Mental Illness (2009); U.S. Department of Veterans Affairs



12



## Suicide > Homicide or War-Related Deaths



Public domain image courtesy of Wikipedia

Every year...

Globally

Almost 1 million suicides

16 per 100,000

1 every 40 seconds

Nationally

Steady rise since 2000

12.3 per 100,000


1 every 13.3 minutes

World Health Organization (2013); American Foundation for Suicide Prevention; Reza et al. (2001)



13





### DoD Suicides & Suicide Rates by Service: Active Component

	All Services	Air Force	Army	Marine Corps	Navy	General Population (CY 2013)
<b>Total Count</b>	259	48	123	45	43	41,149
<b>Rate/100K</b>	18.7	14.4	23.0	23.1	13.4	13.0

Smolenski et al. (2014); Centers for Disease Control and Prevention (2014)

### DoD Suicides & Suicide Rates by Service: Selected Reserve

	All Reserve	Air Force Reserve	Army Reserve	Marine Corps Reserve	Navy Reserve	All National Guard	Air National Guard	Army National Guard
<b>Total Count</b>	87	12	60	11	4	133	14	119
<b>Rate/100K</b>	23.4	--	30.1	--	--	28.9	--	33.4

Smolenski et al. (2014)

### Veteran Suicide Rates

Approx 22% of US suicides each year are veterans

On average, 22 veterans die by suicide each day

U.S. Army photo by Adam Skoczylas



## Veteran Suicide Rates

Male and female veterans had higher firearm suicide rates than nonveterans



DoD photo by Sgt. Michael J. MacLeod, U.S. Army

Kaplan et al (2009)



18



## Etiology of Depression and Suicide




19



## Why Do Some Service Members Develop Depression?

- Physiological
  - Genetics
  - Biological factors
  - Substance abuse
- Psychological
  - Learned helplessness/hopelessness
  - Cognitive factors/Irrational thought processes
- Environmental
  - Loss of loved one
  - Social withdrawal
  - Stress



U.S. Army photo by Ian Graham



20



## Warriors See the World Differently



Photo: Afterdeployment.org



21



## Mental Health Culture vs. Military Culture

### Traditional MH Culture

- Individualistic; 1-on-1 approach
- Emotional vulnerability
- Treatment is delivered individually
- Assumes deficiencies/illness
- Symptoms & risk factors

### Military Culture

- Collectivist; in-group identity
- Emotional toughness
- Leaving group for help jeopardizes safety
- Assumes elitism/strength
- Warrior skills & assets

Bryan (2010)



## Military Myths about Depression

I don't need help because ...

- Only weak people get depression
- My depression will go away if I wait it out
- Treatment does not work

If I seek help ...

- Everyone in my unit will know
- I will lose the trust of my unit
- I will lose my leadership role
- I will lose my security clearance
- My career will be hurt
- I will be administratively/medically separated



U.S. Marine photo. No photographer cited.


Defense Centers of Excellence for Psychological Health and Traumatic Brain Injury



## Why Do Some People (including Service Members and Veterans) Die by Suicide?



## Interpersonal-Psychological Theory of Suicide Risk



Joiner (2005)



## 2 Most Significant Contributors to Suicidal Ideation

- ✓ Thwarted belongingness
- ✓ Perceived ineffectiveness/burden




Image courtesy of S. Miquez, Wikipedia Commons license

Joiner (2005)

hjf



26



## Thwarted Belongingness

Need:

1. Frequent interaction w/ others
  2. Persistent feeling of being cared about
- Interactions must be frequent and positive

Joiner (2005)

hjf



27



## Perceived Ineffectiveness/Burden

Feeling ineffective, plus the sense that loved ones are threatened or burdened by this ineffectiveness.

Joiner (2005)

hjf



28



## Acquired Ability

Reduction of fear through repeated self-injury is necessary for serious suicidal behavior to occur (can occur in the short term or over the long term)

1. Previous suicidal behavior
2. Any experience that reduces fear of injury



Public domain images from DEA

Joiner (2005)

hjf




30




## Empirical Support for IPT in Military Populations

- Suicide note communication
  - Hopelessness
  - Perceived burdensomeness



- Verbally and through suicide note
  - Thwarted belongingness



Cox et al. (2011)



## Depressive Spectrum Disorders: Diagnostic Criteria



## DSM-5: Spectrum of Depressive Disorders

Major Depressive Disorder

Persistent Depressive Disorder (Dysthymia)

Premenstrual Dysphoric Disorder

Substance/ Medication-Induced Depressive Disorder

Depressive Disorder Due to Another Medical Condition

Other Specified Depressive Disorder/ Unspecified Depressive Disorder

American Psychiatric Association (2013)



## DSM-5: Major Depressive Episode

**5 or more of the following for a 2-week period (at least one\*):**

- (1) depressed mood most of the day\*
- (2) markedly diminished interest or pleasure in all, or almost all, activities most of the day\*
- (3) significant weight loss/gain or decrease or increase in appetite
- (4) insomnia or hypersomnia nearly every day
- (5) psychomotor agitation or retardation nearly every day
- (6) fatigue or loss of energy nearly every day
- (7) feelings of worthlessness or excessive or inappropriate guilt
- (8) diminished ability to think or concentrate, or indecisiveness
- (9) recurrent thoughts of death, suicidal ideation



American Psychiatric Association (2013)



## Adjustment Disorder with Depressed Mood

- In DSM-5, this diagnosis falls under Trauma-and Stressor-Related Disorders not Depressive Disorders
- If an individual has symptoms meeting criteria for a major depressive disorder in response to a stressor, the diagnosis of adjustment disorder does not apply



## Trauma and Depression

- Trauma reactions do not *only* include PTSD
- PTSD and depression symptoms overlap, and co-morbidity rates are high
- Some military personnel join the service with a history of depression and/or trauma
- Depression may develop that is not related to deployment or a traumatic event




## Nomenclature For Suicidal And Related Behaviors



## SDV Terminology

- Self-Directed Violence (SDV) Classification System
  - Collaborative approach between the Centers for Disease Control and the VISN 19 MIRECC
  - Describes *thoughts* and *behaviors* associated with suicidality
  - Modifiers exist to address the following:
    - Intent (with, without, or undetermined)
    - Injury (with, without, or fatal)
    - Interrupted act (by self or others)



## Self-Directed Violence Nomenclature: Thoughts

Type	Definition	Modifiers
<b>Non-Suicidal Self-Directed Violence Ideation</b>	Self-reported thoughts regarding a person's desire to engage in self-inflicted potentially injurious behavior. There is no evidence of suicidal intent.	None
<b>Suicidal Ideation</b>	Thoughts of engaging in suicide-related behaviors.	<ul style="list-style-type: none"> <li>▪ Suicidal Intent:               <ul style="list-style-type: none"> <li>- Without</li> <li>- Undetermined</li> <li>- With</li> </ul> </li> </ul>

hjf 40

Brenner et al. (2011)

## Self-Directed Violence Nomenclature: Behaviors

Type	Definition	Modifiers
<b>Preparatory</b>	Acts or preparation towards engaging in Self-Directed Violence, but before potential for injury has begun. This can include anything beyond a verbalization or thought, such as assembling a method or preparing for one's death by suicide.	<ul style="list-style-type: none"> <li>• Suicidal Intent:               <ul style="list-style-type: none"> <li>-Without</li> <li>-Undetermined</li> <li>-With</li> </ul> </li> </ul>
<b>Non-Suicidal Self-Directed Violence</b>	Behavior that is self-directed and deliberately results in injury or the potential for injury to oneself. There is no evidence, whether implicit or explicit, of suicidal intent.	<ul style="list-style-type: none"> <li>• Injury:               <ul style="list-style-type: none"> <li>-Without</li> <li>-With</li> <li>-Fatal</li> </ul> </li> <li>• Interrupted by Self or Other</li> </ul>

hjf 41

Brenner et al. (2011)

## Self-Directed Violence Nomenclature: Behaviors


Type	Definition	Modifiers
<b>Undetermined Self-Directed Violence</b>	Behavior that is self-directed and deliberately results in injury or the potential for injury to oneself. Suicidal intent is unclear based upon the available evidence.	<ul style="list-style-type: none"> <li>• Injury:               <ul style="list-style-type: none"> <li>-Without</li> <li>-With</li> <li>-Fatal</li> </ul> </li> <li>• Interrupted by Self or Other</li> </ul>
<b>Suicidal Self-Directed Violence</b>	Behavior that is self-directed and deliberately results in injury or the potential for injury to oneself. There is evidence, whether implicit or explicit, of suicidal intent.	<ul style="list-style-type: none"> <li>• Injury:               <ul style="list-style-type: none"> <li>-Without</li> <li>-With</li> <li>-Fatal</li> </ul> </li> <li>• Interrupted by Self or Other</li> </ul>

hjf 42

Brenner et al. (2011)

**CDP**

**Suicide Prevention**

**hjf**  43

**CDP** **Shoulder to Shoulder: Finding Strength and Hope Together**



**hjf**  44

Army Suicide Prevention Program (2011)

**CDP** **Goal of Suicide Prevention and Treatment**



**Protective Factors**

**Risk Factors**


**hjf**  48

**CDP**

**Risk Factors, Warning Signs & Protective Factors**

**hjf**  49





**CDP** **Suicide Warning Signs**

- **I** – Ideation
- **S** – Substance Abuse
- **P** – Purposelessness
- **A** – Anxiety
- **T** – Trapped
- **H** – Hopelessness
- **W** – Withdrawal
- **A** – Anger
- **R** – Recklessness
- **M** – Mood Changes

DoD photo by Staff Sgt. Trey Harvey, U.S. Army released.

See handout: "How do you Remember the Warning Signs of Suicide"

Rudd et al. (2006c); American Association of Suicidology (2012)

hjf 51

**CDP** **Suicide Risk Factors**

- More distal in nature than warning signs
- More static in nature than warning signs
- Some risk factors are modifiable/some are not
  - See handout: "Risk Factors for Suicide and Suicidal Behaviors"

American Association of Suicidology (2012)

hjf 52

**CDP** **Military Suicide Risk Factors**

- Relationship problems
- Hopelessness/worthlessness
- Alcohol abuse/dependence
- Feelings of disgrace/isolation
- Guilt or shame
- Stressful military life events
- Easy access to firearms
- Unexplained mood change/depression
- Financial, legal or job performance problems
- Medical or administrative discharge processing
- Sleep problems
- Previous suicide attempts \*\*

Martin et al. (2009); Jones et al. (2012); Ribeiro et al. (2012); Bryan et al. (2013)

hjf 53



## Sleep: Active Duty

Sleep problems outperformed depression and hopelessness as predictors of suicidal ideation and behavior in young adults in the military



U.S. Marine Corps photo by Cpl. Alejandro Pena.

Ribeiro et al. (2012)



## Mental Health Diagnoses and Treatment History: Veterans

Top mental health diagnostic contributors to suicide risk among VA patients:

1. Bipolar disorder
2. Substance use disorders
3. Depression
4. Anxiety disorders other than PTSD

Ilggen et al. (2012)



## TBIs: Active Duty



Multiple TBIs were associated with a significantly higher risk for suicide, even after controlling for symptom severity

Bryan & Clemens (2013)



## Traumatic Brain Injury (TBI): Veterans

- VA patients w/ TBI history 1.55 times more likely to die by suicide than those without
- Among psychiatric inpatients with TBI histories, 27.3% had made a total of 14 suicide attempts



U.S. Army photo. Photographer not listed.

Brenner et al. (2011); Gutierrez et al. (2008)



## Chronic Pain: Veterans



\* Increased risk for suicidal ideation and suicide attempts has been found in individuals with chronic pain, particularly head pain and pain classified as “other non-arthritic”

Juurlink et al. (2004); Fishbain et al. (2009); Ilgen et al. (2008)



## Military Suicide Protective Factors

- Social support or sense of belonging
- Leadership responsibilities
- Effective coping and problem-solving
- Unit cohesion
- Access to assistance services
- Healthy lifestyle promotion
- Spiritual support
- Policies/culture that encourage help-seeking


Martin et al. (2009); Jones et al. (2012); Bryan & Hernandez (2013)





## Assessment of Depression and Suicide






## What Do Depressed or Suicidal Service Members Look Like?






 **Case Study:  
Assessing Mood**



  62


 **Case Study:  
Assessing Other Depressive Symptoms**



  63

 **What Depression and Suicide  
Assessment Tools Do You Use?**



  64

 **Self-Report Screens/ Measures  
of Depressive Symptoms**

**Incorporate self-report measures:**

- PHQ-2 Symptom Checklist = 2 items
- PHQ-9 Symptom Checklist = 9 items
- Center for Epidemiological Studies (CES-D) = 20 items
- Beck Depression Inventory-2 (BDI-2) = 21 items
- Zung Depression Scale = 20 items
- Hamilton Depression Rating Scale = 17 to 31 items

Management of Major Depressive Disorder Working Group (2009); Beck et al. (1996); Carroll et al. (1973); Hamilton (1980); Radloff (1977); Zung (1965)

  65



## Self-Report Measures of Suicidal Ideation/Behavior

- Beck Scale for Suicidal Ideation
  - BSS; Beck & Steer (1991)
- Suicide Intent Scale
  - SIS; Beck, Schuyler, & Herman (1974a)
- Beck Hopelessness Scale
  - BHS; Beck et al. (1974b)
- Suicidal Behavior Questionnaire-18
  - SBQ-18; Linehan (1996)
- Suicidal Behavior Questionnaire-Revised
  - SBQ-R; Osman et al. (2001)



Beck, Schuyler, & Herman (1974a); Beck, et al. (1974b); Beck & Steer (1991); Linehan (1996); Osman, et al. (2001)



## Suicide Risk Assessment

- Previous suicidal behavior
- Current suicidal thoughts, intent, and behavior
- Precipitant stressors (acute and chronic)
- General psychiatric symptoms
- Impulsivity and self-control
- Risk and protective factors
- Use of medications or substances
- Hopelessness
- Warning signs
- Access to lethal means

Department of Veterans Affairs/Department of Defense (2013); Rudd (2006)



## Suicide Risk Assessment (cont.)

- Gain a complete understanding of medical, social and mental health history
- Utilize empirically supported suicide risk assessment instruments in conjunction with a clinical interview
- Obtain collateral information from family, friends, unit, commander, and medical
- Use a direct/nonjudgmental/collaborative approach

*\*Assess risk on an ongoing basis*

Department of Veterans Affairs/Department of Defense (2013)



## Fluid Vulnerability Theory

- Views suicide risk on a continuum
- Acute risk vs. chronic risk
  - Baseline risk - based on personal history, static factors
  - Acute risk - superimposed upon baseline risk
- Suicidal episodes are time limited
- Acute risk resolved when risk factors are effectively targeted

Rudd (2006b)

**CDP**

## Assessing Risk Through Clinical Interview

Assess Baseline Risk

Assess for chronic risk

- Present or absent based on history of multiple attempts

Based on personal history and stable factors

- For example, history of abuse, history of attempts, psychiatric diagnosis

Assess Acute Risk

Reflects the current crisis and overall risk



Exists on a continuum

Time-limited periods of heightened vulnerability to suicide

Includes dynamic factors

- Nature of suicidal thinking, intent, and symptom presentation

Will fluctuate in severity as the suicidal crisis resolves

Rudd (2006a)   70

**CDP**

## Assessing Risk Continued



Acute Risk – Points to Remember

Being thorough does not take a lot of time

Use precise terminology

- Differentiate between non-suicidal thoughts of death, non-suicidal SDV, and suicidal ideation:
  - Non suicidal thoughts of death
  - Non-suicidal SDV
  - Suicidal ideation



“You said that you have had suicidal thoughts. Would you tell me specifically what you’ve been thinking when you think of suicide?”

Rudd (2006a)   71

**CDP**

## Multiple Attempters: Risk Assessment



- Increased vulnerability
- Lower threshold of activation of suicidality
- Always deemed to be at *chronic risk*
- Overall risk level: Always at least “*moderate*” acute/overall risk

Rudd (2006a); Rudd et al. (1996)   72

**CDP**

## Acute Suicide Risk Continuum

Mild	Suicidal ideation of limited frequency, intensity, duration, and specificity. Morbid ideations may be present. There are no identifiable plans; no associated intent; mild dysphoria and related symptoms; good self-control; few other risk factors; and the presence of identifiable protective factors, including social support.
Moderate	Frequent suicidal ideation with limited intensity and duration, some specificity in terms of plans; no intent; good self-control; limited dysphoria and other symptoms; some risk and protective factors, including social support.
High	Frequent, intense, and enduring suicidal ideation; specific plans; some objective markers of intent (e.g., lethal and available method choices, some preparatory behavior); subjective intent may or may not be present; some impairment in self-control; severe dysphoria and/or other symptoms; multiple risk factors present and few protective factors, particularly social support.

Rudd (per discussion with CDP; 2013)   73



## Case Study: Assessing Suicide Risk



## Treatments for Depression



## MDD Psychotherapies

### Efficacious and Specific

- Cognitive Behavior Therapy (CBT)
- Behavior Therapy
- Interpersonal Psychotherapy (IPT)

#### Possibly Efficacious

- Brief Dynamic Therapy
- Emotion-Focused Therapy






U.S. Army/Army Reserve/Army Reserve/Army Reserve



## CBT for Depression: Data from a Meta-Analysis

- Studied in over 75 clinical trials since 1977
- Superior in comparison to waiting list or placebo controls
- No difference in comparison to Behavior Therapy
- Modestly superior in comparison to other therapies
- Significantly better than anti-depressant medication
- Associated with a “preventative” effect







## Cognitive Therapy

### Treatment Approach

- Identify, evaluate, and modify underlying assumptions/ dysfunctional beliefs
- Learn adaptive coping skills
- Break down large problems in smaller steps
- Decision-making via cost-benefit analysis
- Activity scheduling, self-monitoring of mastery and pleasure, and graded task assignments are often used early in therapy



Beck et al (1979); Butler & Beck (1995)

hjf



83



## Behavioral Experiments

*Behavioral experiments can modify a patient's negative beliefs more powerfully than verbal techniques.*

- Designed collaboratively
- Occur during therapy & between sessions
- Goal = an experience that disconfirms the validity of a cognition



Beck (2011)

hjf



84



## General Session Structure

- Mood check
- Bridge from previous session
- Agenda setting
- Review of homework
- Discussion of agenda items
- Periodic summaries
- Homework assignment
- Final summary & feedback



Created by Dave Sattler for the MFRFT program

Wenzel et al. (2011); Photo: Military Family Research Institute, Purdue University

hjf



85



## Behavioral Theory of Depression

- Behavioral patterns associated with depression:
  - Low rate of response-contingent positive reinforcement
  - High rate of punishment
- Central tenet: Depressed individuals do not get enough positive reinforcement from their interactions with the environment to maintain adaptive behavior

Lewinsohn et al. (1980); Wenzel et al. (2011)


hjf



86



## Behavioral Theory of Depression: A Vicious Cycle



Adapted from Lewinsohn et al. (1986)



87



## Behavior Therapy: Behavioral Activation

- Increase pleasurable and mastery activities
- Increase social activities
- Training in social skills, assertiveness, and problem-solving
- Relaxation training and visual imagery
- Behavioral rehearsal and role playing
- Military considerations
  - Exercise may have at one time been pleasurable, but now may be seen as a mastery activity due to mandatory PT/fitness tests
  - May have decreased activity level due to avoidance related to PTSD



88



## Behavior Therapy: Problem-Solving Therapy

- Therapist and client collaboratively identify and prioritize problems, break problems down into manageable tasks, solve problems, and identify coping skills
- Discrete, time-limited, structured intervention



Nezu, Nezu, & Perri (1989)




89



## Interpersonal Psychotherapy (IPT)

- **Goal:** To change behavior by fostering adaptation to current interpersonal roles and situations
- Roots in psychodynamic therapy
- But also draws upon
  - Attachment Theory
  - Increased focus on interpersonal relationship
- More structured than dynamic therapy, but less structured than CBT or BT



U.S. Army photo by Staff Sgt. Helen Miller

Klerman et al (1984)



90



## Treatments for Suicidal Ideation and Behavior



94



## VA/DoD Clinical Practice Guidelines

- Suicide-focused psychotherapy to address suicide risk
  - Clinical Practice Guideline Recommendations:
    - Cognitive Therapy is recommended for non-psychotic patients who survived a recent attempt
    - Problem-solving therapy is recommended for nonpsychotic patients with more than one attempt
- Early evidence-based interventions to target specific symptoms
- Follow-up and monitoring

Department of Veterans Affairs/Department of Defense (2013)



95



## Empirically Supported Treatments/ Interventions

- Dialectical Behavior Therapy (DBT)  
Linehan (1993)
- Means Restriction (Public Health Approach)
  - Hawton (2002), Beautrais (2007), Wiedenmann & Weyerer (1993),  
– Mott et al (2002), Ohberg et al (1995), Law et al (2009)
- Cognitive Therapy for Suicide
  - Brown et al (2005)



96



## Dialectical Behavior Therapy (DBT)

- Goals of DBT according to Linehan:
  - Increase client's behavioral capabilities
  - Improve motivation for skillful behavior through contingency management and reduction of interfering emotions and cognitions
  - Assure generalization of gains to client's environment
  - Structure the treatment environment to reinforce functional rather than dysfunctional behaviors
  - Enhance therapist capabilities and motivation to treat clients effectively

Linehan (1993)



97



## Means Restriction

- Toxic substances
- Medications
- Firearms



## Promising Means Restriction Intervention

- **Means Restriction**
  - Actual process of limiting/removing access to lethal means
- **Means Restriction Counseling**
  - Educate patients and supportive others about risk associated with easy availability of means
  - Collaboratively work with patients and support person to limit/remove access to means until the suicidal risk has lessened



## Means Restriction

Possible mechanisms of effectiveness:

1. Limiting access
2. Reducing opportunity for habituation to fear associated with means for suicide



## Cognitive Therapy for Suicide

Brown et al (2005)



## Results of CT Study

- Significantly fewer suicide attempts in the CT group
- Significantly lower rates of depression in the CT group at 6, 12, and 18 month follow-up
- Significantly lower hopelessness in the CT group at the 6 month point but hopelessness improved overall
- Suicidal ideation went down across the follow-up period but no significant differences between the groups

Brown et al (2005)



## Session #: 1 2 3 4 5 6 7 8 9 10 Early Sessions

- Informed consent
- Treatment engagement
- Assessing level of risk
- Developing a safety plan
- Instilling hope
- Developing a cognitive case conceptualization
- Treatment planning

Wenzel et al (2009)



## Safety Plan vs Safety Contract?



Wenzel, et al. (2009)



## Session #: 1 2 3 4 5 6 7 8 9 10 Middle Sessions

- Modify negative suicide-relevant automatic thoughts & core beliefs
- Teach problem-solving skills
- Help patients develop healthy behavioral coping skills
- Affective coping strategies

Wenzel et al (2009)



## Session #: 1 2 3 4 5 6 7 8 9 10 Middle Sessions

- Identify reasons for living
  - Review advantages and disadvantages of living
- Construct survival kit or hope box
  - Memory aid at time of crisis
    - Photographs
    - Letters
    - Safety plan



Wenzel et al (2009)

hjf



106



## Session #: 1 2 3 4 5 6 7 8 9 10 Middle Sessions

- Build additional coping skills
  - Exercise regimen, hobbies
- Address impulsivity – “procrastinate” suicide
  - Delay tactics
- Increase adaptive use of social support
- Improve compliance w/ adjunctive medical & psychiatric services

Wenzel et al (2009)

hjf



107



## Session #: 1 2 3 4 5 6 7 8 9 10 Later Sessions

- Relapse prevention task
  - Two guided imagery exercises involving past suicidal crisis
  - One guided imagery exercise involving future suicidal crisis
- Debriefing and follow-up
- Additional treatment planning
  - Continuation of treatment
  - Appropriate referrals
  - Termination

Wenzel et al (2009)

hjf



108



## Promising Treatments/ Interventions

- Collaborative Assessment and Management of Suicidality
  - Jobes (2006)
- SAFE Vet
  - Knox et al (2012)
- Means Restriction Counseling
  - Bryan et al (2011)

hjf



109



## Means Restriction Counseling

1. Describe the rationale for means restriction
2. Conduct means restriction counseling
3. Implement the result of Step 2

Rudd & Bryan(2011)



## CDP Website: Deploymentpsych.org

### Features include:

- Descriptions and schedules of upcoming training events
- Blog updated daily with a range of relevant content
- Articles by subject matter experts related to deployment psychology, including PTSD, mTBI, depression, and insomnia
- Other resources and information for behavioral health providers
- Links to CDP's Facebook page and Twitter feed



## Online Learning

The following online courses are located on the CDP website at:

<http://www.deploymentpsych.org/content/online-courses>

**NOTE: All of these courses can be take for free or for CE Credits for a fee**

- Cognitive Processing Therapy (CPT) for PTSD in Veterans and Military Personnel (1.25 CE Credits)
- Prolonged Exposure Therapy for PTSD in Veterans and Military Personnel (1.25 CE Credits)
- Epidemiology of PTSD in Veterans: Working with Service Members and Veterans with PTSD (1.5 CE Credits)
- Provider Resiliency and Self-Care: An Ethical Issue (1 CE Credit)
- Military Cultural Competence (1.25 CE Credits)
- The Impact of Deployment and Combat Stress on Families and Children, Part 1 (2.25 CE Credits)
- The Impact of Deployment and Combat Stress on Families and Children, Part 2 (1.75 CE Credits)
- The Fundamentals of Traumatic Brain Injury (TBI) (1.5 CE Credits)
- Identification, Prevention, & Treatment of Suicidal Behavior in Service Members & Veterans (2.25 CE Credits)
- Depression in Service Members and Veterans (1.25 CE Credits)

*All of these courses and several others are contained in the Serving Our Veterans Behavioral Health Certificate program, which also includes 20+ hours of Continuing Education Credits for \$350.*



## Provider Support

CDP's "Provider Portal" is exclusively for individuals trained by the CDP in evidence-based psychotherapies (e.g., CPT, PE, and CBT-I)

### Features include:

- Consultation message boards
- Hosted consultation calls
- Printable fact sheets, manuals, handouts, and other materials
- FAQs and one-on-one interaction with answers from SMEs
- Videos, webinars, and other multimedia training aids

Participants in CDP's evidence-based training will automatically receive an email instructing them how to activate their user name and access the "Provider Portal" section at Deploymentpsych.org.





## How to Contact Us

### Center for Deployment Psychology

Department of Medical & Clinical Psychology  
Uniformed Services University of the Health Sciences  
4301 Jones Bridge Road, Executive Office: Bldg. 11300-602  
Bethesda, MD 20813-4768

**Email:** [General@DeploymentPsych.org](mailto:General@DeploymentPsych.org)

**Website:** [DeploymentPsych.org](http://DeploymentPsych.org)

**Facebook:** <http://www.facebook.com/DeploymentPsych>

**Twitter:** @DeploymentPsych





## **Assessment and Treatment of Depression and Suicidal Behavior Associated with Military Service: An Overview**

### **References**

- Abramson, L. Y., Seligman, M. E., Teasdale, J. D. (1978). Learned helplessness in humans; critique and reformulation. *Journal of Abnormal Psychology*, (87), 1, 49-74.
- Adler, A.B, McGurk, D.M, Stetz, MC & Bliese, P.D. (2003). *Military occupational stressors in garrison, training, & deployed environments. Work, stress, & health: New challenges for a changing workplace.* Conference sponsored by the National Institute of Occupational Safety & Health and the American Psychological Association, Toronto, Canada. Retrieved from: <http://www.dtic.mil/cgi-bin/GetTRDoc?AD=ada425834>
- American Association of Suicidology (2012). *Risk Factors for Suicide and Suicidal Behaviors.* American Association of Suicidology, Washington DC. Retrieved from: [http://www.suicidology.org/Portals/14/docs/Resources/FactSheets/Risk-Factors\\_2013.pdf](http://www.suicidology.org/Portals/14/docs/Resources/FactSheets/Risk-Factors_2013.pdf)
- American Foundation for Suicide Prevention (n.d.). *Facts and Figures.* Retrieved from: <http://www.afsp.org/understanding-suicide/facts-and-figures>
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC
- Army Suicide Prevention Program. (2011). *Shoulder to Shoulder: Finding Hope and Strength Together.* Retrieved from: <http://www.armyg1.army.mil/hr/suicide/media2asp>.
- Beautrais, A. (2007). Suicide by jumping: a review of research and prevention strategies. *Crisis*, 28 (Suppl 1), 58- 63.
- Beck, A.T. (1976). *Cognitive therapies and emotional disorders.* New York: New American Library.
- Beck, A.T., Rush, A.J., Shaw, B.F., & Emery, G. (1979). *Cognitive therapy of depression.* New York: Guilford.
- Beck, A. T., Schuyler, D., & Herman, I. (1974a). Development of suicidal intent scales. In A. T. Beck, H. Resnik, & D. J. Lettieri (Eds.), *The prediction of suicide.* Bowie, MD: Charles Press.
- Beck, A.T., & Steer, R.A. (1991). *Manual for the Beck scale for suicide ideation.* San Antonio, TX: Psychological Corp.
- Beck, A.T., Steer, R.A. & Brown, G.K. (1996). *Manual for the Beck Depression Inventory-II.* San Antonio, TX: Psychological Corp.

- Beck, A. T., Weissman, A., Lester, D., & Trexler, L. (1974b). The measurement of pessimism: The Hopelessness Scale. *Journal of Consulting and Clinical Psychology*, 42(6), 861-865.
- Beck, J. S. (2011). *Cognitive behavior therapy: Basics and beyond* (2<sup>nd</sup> ed.). NY: Guilford Press.
- Brenner, L.A., Breshears, R.E., Betthausen, L.M., Bellon, K.K., Holman, E., Harwood, J.E.F., et al. (2011). Implementation of a suicide nomenclature within two VA healthcare settings. *Journal of Clinical Psychology in Medical Settings*, 18(2), 116-128.
- Brenner, L.A., Ignacio, R.V. Y Blow, F.C. (2011). Suicide and traumatic brain injury among individuals seeking Veterans Health Administration services. *The Journal of Head Trauma Rehabilitation*, 26 (4), 257- 264.
- Brown, G. K., Ten Have, T., Henriques, G. R., Xie, S. X., Hollander, J. E., & Beck, A. T. (2005). Cognitive therapy for the prevention of suicide attempts: A randomized controlled trial. *Journal of the American Medical Association*, 294(5), 563-570.
- Bush, N. E., Dobscha, S. K., Crumpton, R., Denneson, L. M., Hoffman, J. E., Crain, A., Cromer, R., and Kinn, J. T. (2014). A virtual hope box smartphone app as an accessory to therapy: proof of concept in a clinical sample of veterans. *Suicide and Life Threatening Behavior*, 1-9.
- Bryan, C. J. (2010). *Military culture, mental health stigma, and new approaches to mental health service delivery* [PowerPoint slides]. Retrieved from [http://www.health.mil/dhb/downloads/TaskForce2010/6\\_March%2011%20Task%20Force%20Meeting%20Materials/Presentations/Bryan\\_DOD%20Suicide%20Task%20Force%20presentation.pdf](http://www.health.mil/dhb/downloads/TaskForce2010/6_March%2011%20Task%20Force%20Meeting%20Materials/Presentations/Bryan_DOD%20Suicide%20Task%20Force%20presentation.pdf)
- Bryan, C., Bryan, A. O., Ray-Sannerud, B. N., Etienne, N., and Morrow, C. (2013). Suicide attempts before joining the military increase risk for suicide attempts and severity of suicidal ideation among military personnel and veterans. *Comprehensive Psychiatry*.
- Bryan, C. and Clemans, T. (2013). Repetitive traumatic brain injury, psychological symptoms, and suicide risk in a clinical sample of deployed military personnel. *Journal of the American Medical Association, Psychiatry*, 70 (7), 686-691.
- Bryan, C. and Hernandez, A. M. (2013). The functions of social support as protective factors for suicidal ideation in a sample of Air Force personnel. *Suicide and Life-Threatening Behavior* 43 (5). The American Association of Suicidology.
- Bryan, C., Morrow, C., Etienne, N., Ray-Sannerud, B. (2013). Guilt, shame, and suicidal ideation in a military outpatient clinical sample. *Depression and Anxiety*, 30, 55-60.
- Bryan, C. J., Stone, S. L., & Rudd, D. M. (2011). A practical, evidence-based approach for means-restriction counseling with suicidal patients. *Professional Psychology, Research and Practice*, 42(5), 339-346.
- Burns, D. D. (1980). *Feeling good: The new mood therapy*. New York: New American Library.

- Butler, A.C. & Beck, A.T. (1995). Cognitive therapy for depression. *The Clinical Psychologist*, 48(3), 3-5.
- Butler, A.C., Chapman, J.E., Forman, E.M., & Beck, A.T. (2006). The empirical status of cognitive-behavioral therapy: A review of meta-analyses. *Clinical Psychology Review*, 26, 17-31.
- Carroll, B.J., Fielding, J.M., & Blashki, T.G. (1973). Depression rating scales. *Archives of General Psychiatry*, 28, 361-366.
- Centers for Disease Control and Prevention. (2012). *National Suicide Statistics at a Glance*. Retrieved from <http://www.cdc.gov/violenceprevention/suicide/statistics/index.html>.
- Conner, K. R., & Zhong, Y. (2003). State firearm laws and rates of suicide in men and women. *American journal of preventive medicine*, 25(4), 320-324.
- Conwell, Y., Duberstein, P. R., & Caine, E. D. (2002). Risk factors for suicide in later life. *Biological psychiatry*, 52(3), 193-204.
- Cornette, M. (2013). *Conceptual model of suicide risk*.
- Cox, D. W., Ghahramanlou-Holloway, M., Greene, F., Bakalar, J., Schendel, C., Nademin, M. E., Jobes, D. A., Englert, D. R., & Kindt, M. (2011). Suicide in the United States Air Force: Risk factors communicated before and at death. *Journal of Affective Disorders*, 133, 398-405.
- Crosby, A.E., Ortega, L., & Melanson C. (2011). *Self-directed violence surveillance: Uniform definitions and recommended data elements, version 1.0*. Atlanta, GA: Centers for Disease Control and Prevention , National Center for Injury Prevention and Control.
- Defense Centers of Excellence for Psychological Health and Traumatic Brain Injury (n.d.). *Dispelling the Myths About Depression*. Retrieved from: <http://www.dcoe.mil/content/navigation/documents/Dispelling%20the%20Myths%20About%20Depression.pdf>
- Department of Defense. (2013). *2011 health related behaviors survey of active duty military personnel*. Retrieved from: <http://Tricare.mil/tma/dhcape/surveys/coresurveys/surveyhealthrelatedbehaviors/downloads/Final%202011%20HRB%20Active%20Duty%20Survey%20Summary.pdf>
- Department of Veterans Affairs/Department of Defense (2013). *VA/DOD clinical practice guideline for assessment and management of patients at risk for suicide*. Washington, D. C.
- Department of Veterans Affairs/Department of Defense (2013). *VA/DoD CPG for Assessment and Management of Patients at Risk for Suicide*. Washington DC.
- Fishbain, D.A., Bruns, D., Disorbio, J.M. & Lewis, J.E. (2009). Risk for five forms of suicidality

- in acute pain patients and chronic pain patients vs pain-free community controls. *Pain Medicine*, 10 (6), 1095- 1105
- Gloaguen, V., Cottraux, J., Cucherat, M., & Blackburn, I. (1998). A meta-analysis of the effects of cognitive therapy in depressed patients. *Journal of Affective Disorders*, 49(1), 59-72.
- Gutierrez, P.M., Brenner, L.A., & Huggins, J.A. (2008). A preliminary investigation of suicidality in psychiatrically hospitalized veterans with traumatic brain injury. *Archives of Suicide Research*, 12, 336- 343.
- Halvorson, A. (2010). *Understanding the military: The institution, the culture, & the people*. Retrieved from: [http://www.samhsa.gov/sites/default/files/military\\_white\\_paper\\_final.pdf](http://www.samhsa.gov/sites/default/files/military_white_paper_final.pdf)
- Hamilton, M. (1980). Rating depressive patients. *Journal of Clinical Psychiatry*, 41, 21-24.
- Hawton, K. (2002). United Kingdom legislation on pack sizes of analgesics: background, rationale, and effects on suicide and deliberate self-harm. *Suicide and Life-Threatening Behavior*, 32, 223-229.
- Hollon, S.D. & Ponniah, K. (2010). A review of empirically supported psychological therapies for mood disorders in adults. *Depression and Anxiety*, 27 (10), 891- 932.
- Ilgén, M. A., McCarthy, J. F., Ignacio, R. V., Bohnert, A. S., Valenstein, M., Blow, F. C. (2012). Psychopathology, Iraq, and Afghanistan service, and suicide among Veterans Health Administration patients. *Journal of Consulting and Clinical Psychology*, 80 (3), 323-330.
- Ilgén, M.A., Zivin, K., McCammon, R.J., and Valenstein, M. (2008). Pain and suicidal thoughts, plans, and attempts in the United States. *General Hospital Psychiatry*, 30 (6), 521-527.
- Jacobson, N.S., Martell, C.R. & Dimidjian, S. (2001). Behavioral Activation for depression: Returning to contextual roots. *Clinical Psychology: Science and Practice*, 8, 255–270.
- Jobes, D.A. (2006). *Managing suicidal risk: a collaborative approach*. New York, NY: The Guilford Press.
- Joiner, T. (2005). *Why people die by suicide*. Cambridge, MA: Harvard University Press.
- Jones, D.E., Hourani, L.L., Rariden, M.B., Hammond, P.J., & Werbel, A.D. (2012). Suicide prevention in the military. In C.H. Kennedy & E. A. Zilmer (Eds). *Military psychology: Clinical and operational applications* (211-250). New York, NY: The Guilford Press.
- Juurlink, D.N., Herrmann, N., & Szalai, J.P. (2004). Medical illness and the risk of suicide in the elderly. *Archives of Internal Medicine*, 164, 1170- 1184.
- Kemp, J., & Bossarte, R. (2013). *Suicide Data Report: 2012*. Department of Veterans Affairs, Mental Health Services, Suicide Prevention Program.

- Kaplan, M.S., McFarland, B.H., & Huguet, N. (2009). Characteristics of adult male and female firearm suicide decedents: findings from the National Violent Death Reporting System. *Injury Prevention, 15* (5), 322-327.
- Klerman, G.L., Weissman, M.M., Rounsaville, B.J. & Chevron, E.S. (1984). *Interpersonal psychotherapy of depression*. Basic Books: New York, New York.
- Knox, K., Stanley, B., Currier, G., Brenner, L., Ghahramanlou-Holloway, M., & Brown, G. (2012). An emergency department-based brief intervention for Veterans at risk for suicide (SAFE VET). *American Journal of Public Health, 102*(S1), S33-S37.
- Kupfer D.J. (1991). Long-term treatment of depression. *Journal of Clinical Psychiatry, 52*, 28-34.
- Law, C.K., Yip, P.S.F., Wincy, S.C.C. et al (2009). Evaluating the effectiveness of barrier installation for preventing railway suicides in Hong Kong. *Journal of Affective Disorders, 114*, 254- 262.
- Lewinsohn, P.M., Munoz, R.F., Youngren, M.A., & Zeiss, A.M. (1986). *Control your depression*. NY: Simon & Schuster.
- Lewinsohn, P.M., Sullivan, J.M., & Grosscup, S.J. (1980). Changing reinforcing events: An approach to the treatment of depression. *Psychotherapy: Theory, Research, & Practice, 17*(3), 322-334.
- Linehan, M.M. (1993). *Cognitive-Behavioral Treatment of Borderline Personality Disorder*. New York, NY: The Guilford Press.
- Linehan, M.M. (1996). *Suicidal Behaviors Questionnaire (SBQ)*. University of Washington, Seattle.
- Luxton, D.D., Osenbach, J.E., Reger, M.A., Smolenski, D.J., Skopp, N.A., Bush, N.E., & Gahm, G.A. (2012). *Department of Defense Suicide Event Report Calendar Year 2011 Annual Report*. National Center for Telehealth & Technology.
- Management of Major Depressive Disorder Working Group (2009). *VA/DoD Clinical Practice Guideline: Management of Major Depressive Disorder*. Washington, DC, Department of Veteran's Affairs, Department of Defense. Retrieved from <http://www.healthquality.va.gov/mdd>
- Martell, C.R., Addis, M.E., & Jacobson, N.S. (2001). *Depression in context: Strategies for guided action*. New York: Norton.
- Martell, C.R., Dimidjian, S., and Herman-Dunn, R. (2010). *Behavioral activation for depression: A clinician's guide*. New York: Guilford Press.

- Martin, J., Ghahramanlou-Holloway, M., Lou, K., Tucciarone, P. (2009). A comparative review of U.S. military and implications for OEF/OIF suicide prevention efforts. *Journal of Mental Health Counseling*, 31(2), 101-118...
- Miles, D. (2010, April 23). VA strives to prevent veteran suicides. *American Forces Press Service*. Retrieved from: <http://www.mirecc.va.gov/visn19/education/nomenclature.asp>.
- Mott, J.A., Wolfe, M.I., Alverson, C.J. et al. (2002). National vehicle emissions policies and practices and declining U.S. carbon monoxide-related mortality. *Journal of the American Medical Association*, 288, 988- 995.
- National Alliance on Mental Illness (2009). *Depression and Veterans Fact Sheet*. Retrieved from: <http://www.nami.org/Template.cfm?Section=Depression&Template=/ContentManagement/ContentDisplay.cfm&ContentID=87942>
- Nezu, A.M. Nezu, C.M. and Perri, M.G. (1989). *Problem-solving therapy for depression: theory, research, and clinical guidelines*. New York: Wiley.
- Ohberg, A., Lonnqvist, J., Sarna, S. et al. (1995). Trends and availability of suicide methods in Finland: proposals for restrictive measures. *British Journal of Psychiatry*, 166, 35- 43.
- O'Reilly, H. (2013). Cognitive model image (tree metaphor).
- Osman, A., Bagge, C.L., Guitierrez, P.M., Konic, L.C., Kooper, B.A., & Barrios, F.X. (2001). The Suicidal Behaviors Questionnaire-Revised (SBQ-R): Validation with clinical and nonclinical samples, *Assessment* 8(4), 443-454.
- Pigeon, W.R., Piquart, M., & Conner, K. (2012). Meta-analysis of sleep disturbance and suicidal thoughts and behaviors. *Journal of Clinical Psychiatry*, 73(9), 1160-1167).
- Radloff, L.S. (1977). The CES-D scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1, 385-401.
- Reza, A., Mercy, J.A., & Krug, E. (2001). Epidemiology of violent deaths in the world. *Injury Prevention*, 7, 104-111
- Ribeiro J. D., Pease J. L., Gutierrez P. M., Silva, C., Bernert, R. A., Rudd, M. D., & Joiner, T. E. (2012). Sleep problems outperform depression and hopelessness as cross-sectional and longitudinal predictors of suicidal ideation and behavior in young adults in the military. *Journal of Affective Disorders*, 136(3), 743-50.
- Rudd, M.D. (2006a). *The assessment and management of suicidality*. Sarasota, FL: Professional Resource Exchange.
- Rudd, M.D. (2006b). Fluid vulnerability theory: A cognitive approach to understanding the process of acute and chronic suicide risk. In T.E. Ellis' (Ed.) *Cognition and suicide: Theory, research, and therapy* (pp. 355-380). Washington, DC: American Psychological Association.

- Rudd, M.D., Berman, A.L., Joiner, T.E., Nock, M.K., Silverman, M.M., Mandrusiak, M., Van Orden, K., & Witte, T. (2006c). Warning signs for suicide: Theory, research, and clinical applications. *Suicide and Life-Threatening Behavior*, 36(3), 255-262.
- Rudd, M.D. & Bryan, C.J. (2011). *Brief cognitive behavioral therapy for suicidal soldiers*. Unpublished manual, 17-21.
- Rudd, M. D., Joiner, T.E., & Rajab, M.H. (2001). *Treating suicidal behavior: An effective, time-limited approach*. New York, NY: Guilford Press.
- Segal, Z.V., Williams, J.M.G., and Teasdale, J.D. (2013). *Mindfulness-based cognitive therapy for depression (2<sup>nd</sup> edition)*. New York: The Guilford Press.
- Smolenski, D.J., Reger, M.A., Alexander, C.L., Skopp, N.A., Bush, N.E., Luxton, D.D., & Gahm, G.A. (2013). *Department of Defense Suicide Event Report Calendar Year 2012 Annual Report*. National Center for Telehealth & Technology Defense Centers of Excellence for Psychological Health & Traumatic Brain Injury.
- Tanielian, T. L., & Jaycox, L. H. (Eds.). (2008). *Invisible wounds of war: Psychological and cognitive injuries, their consequences, and services to assist recovery*. Rand Corporation.
- Thomas, J. L., Wilk, J.E., Riviere, L.A., McGurk, D., Castro, C.A., & Hoge, C.W. (2010). Prevalence of mental health problems and functional impairment among active component and National Guard soldiers 3 and 12 months following combat in Iraq. *Archives of General Psychiatry*, 67(6), 614-623.
- Wampold, B. E., Minami, T., Baskin, T. W. & Tierney, S.C. (2002). A meta-(re) analysis of the effects of cognitive therapy versus 'other therapies' for depression. *Journal of Affective Disorders*, 68, 159-165.
- Wells, T.S., LeardMann, C.A., Fortuna, S.O., Smith, B., Smith, T.C., Ryan, M.A.K, Blazer, D. (2010). A prospective study of depression following combat deployment in support of the wars in Iraq & Afghanistan. *American Journal of Public Health*, 10(1), 90-99.
- Wenzel, A., Brown, G. K., & Beck, A. T. (2009). *Cognitive therapy for suicidal patients*. Washington, DC: American Psychological Association.
- Wenzel, A., Brown, G.K., & Karlin, B.E. (2011). *Cognitive behavioral therapy for depression in veterans and military servicemembers: Therapist manual*. Washington, DC: U.S. Department of Veterans Affairs.
- Weidenmann, A. & Weyerer, S. (1993). The impact of availability, attraction and lethality of suicide methods on suicide rates in Germany. *Acta Psychiatrica Scandinavica*, 88, 364-368
- World Health Organization. (2013). *Suicide prevention*. Retrieved from: [http://www.who.int/mental\\_health/prevention/suicide/suicideprevent/en/](http://www.who.int/mental_health/prevention/suicide/suicideprevent/en/).
- Zung, W.W. (1965). A self-rating depression scale. *Archives of General Psychiatry*, 12, 63-70.





# PHQ-9 — Nine Symptom Checklist

**Patient Name** \_\_\_\_\_ **Date** \_\_\_\_\_

1. Over the last 2 weeks, how often have you been bothered by any of the following problems? Read each item carefully, and circle your response.

a. Little interest or pleasure in doing things

**Not at all**      **Several days**      **More than half the days**      **Nearly every day**

b. Feeling down, depressed, or hopeless

**Not at all**      **Several days**      **More than half the days**      **Nearly every day**

c. Trouble falling asleep, staying asleep, or sleeping too much

**Not at all**      **Several days**      **More than half the days**      **Nearly every day**

d. Feeling tired or having little energy

**Not at all**      **Several days**      **More than half the days**      **Nearly every day**

e. Poor appetite or overeating

**Not at all**      **Several days**      **More than half the days**      **Nearly every day**

f. Feeling bad about yourself, feeling that you are a failure, or feeling that you have let yourself or your family down

**Not at all**      **Several days**      **More than half the days**      **Nearly every day**

g. Trouble concentrating on things such as reading the newspaper or watching television

**Not at all**      **Several days**      **More than half the days**      **Nearly every day**

h. Moving or speaking so slowly that other people could have noticed. Or being so fidgety or restless that you have been moving around a lot more than usual

**Not at all**      **Several days**      **More than half the days**      **Nearly every day**

i. Thinking that you would be better off dead or that you want to hurt yourself in some way

**Not at all**      **Several days**      **More than half the days**      **Nearly every day**

2. If you checked off any problem on this questionnaire so far, how difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?

**Not Difficult at All**      **Somewhat Difficult**      **Very Difficult**      **Extremely Difficult**

# PHQ-9 — Scoring Tally Sheet

Patient Name \_\_\_\_\_ Date \_\_\_\_\_

**1. Over the last 2 weeks, how often have you been bothered by any of the following problems? Read each item carefully, and circle your response.**

	Not at all	Several days	More than half the days	Nearly every day
	0	1	2	3
a. Little interest or pleasure in doing things				
b. Feeling down, depressed, or hopeless				
c. Trouble falling asleep, staying asleep, or sleeping too much				
d. Feeling tired or having little energy				
e. Poor appetite or overeating				
f. Feeling bad about yourself, feeling that you are a failure, or feeling that you have let yourself or your family down				
g. Trouble concentrating on things such as reading the newspaper or watching television				
h. Moving or speaking so slowly that other people could have noticed. Or being so fidgety or restless that you have been moving around a lot more than usual				
i. Thinking that you would be better off dead or that you want to hurt yourself in some way				
Totals				

**2. If you checked off any problem on this questionnaire so far, how difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?**

Not Difficult At All	Somewhat Difficult	Very Difficult	Extremely Difficult
0	1	2	3

# How to Score PHQ-9

## Scoring Method For Diagnosis

### Major Depressive Syndrome is suggested if:

- Of the 9 items, 5 or more are circled as at least "More than half the days"
- Either item 1a or 1b is positive, that is, at least "More than half the days"

### Minor Depressive Syndrome is suggested if:

- Of the 9 items, b, c, or d are circled as at least "More than half the days"
- Either item 1a or 1b is positive, that is, at least "More than half the days"

## Scoring Method For Planning And Monitoring Treatment

### Question One

- To score the first question, tally each response by the number value of each response:

Not at all = 0

Several days = 1

More than half the days = 2

Nearly every day = 3

- Add the numbers together to total the score.
- Interpret the score by using the guide listed below:

Score	Action
$\leq 4$	The score suggests the patient may not need depression treatment.
$> 5-14$	Physician uses clinical judgment about treatment, based on patient's duration of symptoms and functional impairment.
$\geq 15$	Warrants treatment for depression, using antidepressant, psychotherapy and/or a combination of treatment

### Question Two

In question two the patient responses can be one of four: not difficult at all, somewhat difficult, very difficult, extremely difficult. The last two responses suggest that the patient's functionality is impaired. After treatment begins, the functional status is again measured to see if the patient is improving.





## **Risk Factors for Suicide and Suicidal Behaviors I.**

**Chronic Risk Factors** (If present, these increase risk over one's lifetime.)

### **A. Perpetuating Risk Factors – permanent and non-modifiable**

- Demographics: White, American Indian, Male, Older Age (review current rates<sup>1</sup>), Separation or Divorce, Early Widowhood
- History of Suicide Attempts – especially if repeated
- Prior Suicide Ideation
- History of Self-Harm Behavior
- History of Suicide or Suicidal Behavior in Family
- Parental History of:
  - Violence
  - Substance Abuse (Drugs or Alcohol)
  - Hospitalization for Major Psychiatric Disorder
  - Divorce
- History of Trauma or Abuse (Physical or Sexual)
- History of Psychiatric Hospitalization
- History of Frequent Mobility
- History of Violent Behaviors
- History of Impulsive/Reckless Behaviors

### **Predisposing and Potentially Modifiable Risk Factors**

- Major Axis I Psychiatric Disorder, especially:
  - Mood Disorder
  - Anxiety Disorder
  - Schizophrenia
  - Substance Use Disorder (Alcohol Abuse or Drug Abuse/Dependence)
  - Eating Disorders
  - Body Dysmorphic Disorder
  - Conduct Disorder
- Axis II Personality Disorder, especially Cluster B

---

<sup>1</sup> Available from <http://webapp.cdc.gov/sasweb/ncipc/mortrate.html>

- Axis III Medial Disorder, especially if involves functional impairment and/or chronic pain)
- Traumatic Brain Injury
- Co-morbidity of Axis I Disorders (especially depression and alcohol misuse), of Axis I and Axis II (especially if Axis II Disorder is Antisocial PD or Borderline PD), of Axis I and Axis III Disorders
- Low Self-esteem/High Self-hate
- Tolerant/Accepting Attitude Toward Suicide
- Exposure to Another's Death by Suicide
- Lack of Self or Familial Acceptance of Sexual Orientation
- Smoking
- Perfectionism (especially in context of depression)

## **Risk Factors for Suicide and Suicidal Behaviors II**

### **Contributory Risk Factors**

- Firearm Ownership or Easy Accessibility
- Acute or Enduring Unemployment
- Stress (job, marriage, school, relationship...)

### **Acute Risk Factors (If present, these increase risk in the near-term)**

- Demographics: Recently Divorced or Separated with Feelings of Victimization or Rage
- Suicide Ideation (threatened, communicated, planned, or prepared for)
- Current Self-harm Behavior
- Recent Suicide Attempt
- Excessive or Increased Use of Substances (alcohol or drugs)
- Psychological Pain (acute distress in response to loss, defeat, rejection, etc.)
- Recent Discharge from Psychiatric Hospitalization
- Anger, Rage, Seeking Revenge
- Aggressive Behavior
- Withdrawal from Usual Activities, Supports, Interests, School or Work; Isolation (e.g. lives alone)
- Anhedonia
- Anxiety, Panic
- Agitation
  - Insomnia
  - Persistent Nightmares

- Suspiciousness, Paranoia (ideas of persecution or reference)
- Severe Feelings of Confusion or Disorganization
- Command Hallucinations Urging Suicide
- Intense Affect States (e.g. desperation, intolerable aloneness, self-hate...)
- Dramatic Mood Changes
- Hoplessness, Poor Problem-solving, Cognitive Constriction (thinking in black and white terms, not able to see gray areas, alternatives...), Rumination, Few Reasons for Living, Inability to Imagine Possibly Positive Future Events
- Perceived Burdensomeness
- Recent Diagnosis of Terminal Condition
- Feeling Trapped, Like There is No Way Out (other than death); Poor Problem-Solving
- Sense of Purposelessness or Loss of Meaning; No Reasons for Living
- Negative or Mixed Attitude Toward Help-Receiving
- Negative or Mixed Attitude by Potential Caregiver to Individual
- Recklessness or Excessive Risk-Taking Behavior, Especially if Out of Character or Seemingly Without Thinking of Consequences, Tendency Toward Impulsivity

**Precipitating or Triggering Stimuli (Heighten Period of Risk if Vulnerable to Suicide)**

- Any Real or Anticipated Event Causing or Threatening:
  - Shame, Guilt, Despair, Humiliation, Unacceptable Loss of Face or Status
  - Legal Problems (loss of freedom), Financial Problems, Feelings of Rejection/Abandonment
- Recent Exposure to Another's Suicide (of friend or acquaintance, of celebrity through media...)



## **American Association of Suicidology**

AAS is a membership organization for all those involved in suicide prevention and intervention, or touched by suicide. AAS is a leader in the advancement of scientific and programmatic efforts in suicide prevention through research, education and training, the development of standards and resources, and survivor support services. For membership information, please contact:

*American Association of Suicidology*

*5221 Wisconsin Ave., N.W.*

*Second Floor*

*Washington, DC 20015*

*tel. (202) 237-2280*

*fax (202) 237-2282*

*[www.suicidology.org](http://www.suicidology.org)*

*[info@suicidology.org](mailto:info@suicidology.org)*

**If you or someone you know is  
suicidal, please contact a mental  
health professional or call 1-800-  
273-TALK (8255).**

### Self-Directed Violence Classification System\*

Type	Sub-Type	Definition	Modifiers	Terms
Thoughts	Non-Suicidal Self-Directed Violence Ideation	Self-reported thoughts regarding a person's desire to engage in self-inflicted potentially injurious behavior. There is no evidence of suicidal intent.  For example, persons engage in Non-Suicidal Self-Directed Violence Ideation in order to attain some other end (e.g., to seek help, regulate negative mood, punish others, to receive attention).	N/A	•Non-Suicidal Self-Directed Violence Ideation
	Suicidal Ideation	Thoughts of engaging in suicide-related behavior.  For example, intrusive thoughts of suicide without the wish to die would be classified as Suicidal Ideation, Without Intent.	•Suicidal Intent: -Without -Undetermined -With	•Suicidal Ideation, Without Suicidal Intent •Suicidal Ideation, With Undetermined Suicidal Intent •Suicidal Ideation, With Suicidal Intent
Behaviors	Preparatory	Acts or preparation towards engaging in Self-Directed Violence, but before potential for injury has begun. This can include anything beyond a verbalization or thought, such as assembling a method (e.g., buying a gun, collecting pills) or preparing for one's death by suicide (e.g., writing a suicide note, giving things away).  For example, hoarding medication for the purpose of overdosing would be classified as Suicidal Self-Directed Violence, Preparatory.	•Suicidal Intent: -Without -Undetermined -With	•Non-Suicidal Self-Directed Violence, Preparatory •Undetermined Self-Directed Violence, Preparatory •Suicidal Self-Directed Violence, Preparatory
	Non-Suicidal Self-Directed Violence	Behavior that is self-directed and deliberately results in injury or the potential for injury to oneself. There is no evidence, whether implicit or explicit, of suicidal intent.  For example, persons engage in Non-Suicidal Self-Directed Violence in order to attain some other end (e.g., to seek help, regulate negative mood, punish others, to receive attention).	•Injury: -Without -With -Fatal •Interrupted by Self or Other	•Non-Suicidal Self-Directed Violence, Without Injury •Non-Suicidal Self-Directed Violence, Without Injury, Interrupted by Self or Other •Non-Suicidal Self-Directed Violence, With Injury •Non-Suicidal Self-Directed Violence, With Injury, Interrupted by Self or Other •Non-Suicidal Self-Directed Violence, Fatal
	Undetermined Self-Directed Violence	Behavior that is self-directed and deliberately results in injury or the potential for injury to oneself. Suicidal intent is unclear based upon the available evidence.  For example, the person is unable to admit positively to the intent to die (e.g., unconsciousness, incapacitation, intoxication, acute psychosis, disorientation, or death); <b>OR</b> the person is reluctant to admit positively to the intent to die for other or unknown reasons.	•Injury: -Without -With -Fatal •Interrupted by Self or Other	•Undetermined Self-Directed Violence, Without Injury •Undetermined Self-Directed Violence, Without Injury, Interrupted by Self or Other •Undetermined Self-Directed Violence, With Injury •Undetermined Self-Directed Violence, With Injury, Interrupted by Self or Other •Undetermined Self-Directed Violence, Fatal
	Suicidal Self-Directed Violence	Behavior that is self-directed and deliberately results in injury or the potential for injury to oneself. There is evidence, whether implicit or explicit, of suicidal intent.  For example, a person with the wish to die cutting her wrists with a knife would be classified as Suicide Attempt, With Injury.	•Injury: -Without -With -Fatal •Interrupted by Self or Other	•Suicide Attempt, Without Injury •Suicide Attempt, Without Injury, Interrupted by Self or Other •Suicide Attempt, With Injury •Suicide Attempt, With Injury, Interrupted by Self or Other •Suicide

\* Developed in collaboration with the Centers for Disease Control and Prevention

### Self-Directed Violence Classification System\*

<b>Key Terms</b>	<p><b><i>Self-Directed Violence:</i></b> Behavior that is self-directed and deliberately results in injury or the potential for injury to oneself.</p> <p><b><i>Suicidal Intent:</i></b> There is past or present evidence (implicit or explicit) that an individual wishes to die, means to kill him/herself, and understands the probable consequences of his/her actions or potential actions. Suicidal intent can be determined retrospectively and in the absence of suicidal behavior.</p> <p><b><i>Physical Injury:</i></b> A (suspected) bodily lesion resulting from acute overexposure to energy (this can be mechanical, thermal, electrical, chemical, or radiant) interacting with the body in amounts or rates that exceed the threshold of physiological tolerance. In some cases an injury results from an insufficiency of vital elements, such as oxygen. Acute poisonings and toxic effects, including overdoses of substances and wrong substances given or taken in error are included, as are adverse effects and complications of therapeutic, surgical and medical care. Psychological injury is excluded in this context.</p> <p><b><i>Interrupted By Self or Other:</i></b> A person takes steps to injure self but is stopped by self/another person prior to fatal injury. The interruption may occur at any point.</p> <p><b><i>Suicide Attempt:</i></b> A non-fatal self-inflicted potentially injurious behavior with any intent to die as a result of the behavior.</p> <p><b><i>Suicide:</i></b> Death caused by self-inflicted injurious behavior with any intent to die as a result of the behavior.</p>
------------------	---

\* Developed in collaboration with the Centers for Disease Control and Prevention



## How do you Remember the Warning Signs of Suicide?

Here's an Easy-to-Remember Mnemonic:

### ***IS PATH WARM?***

I	Ideation
S	Substance Abuse
P	Purposelessness
A	Anxiety
T	Trapped
H	Hopelessness
W	Withdrawal
A	Anger
R	Recklessness
M	Mood Change

A person in acute risk for suicidal behavior most often will show:

Warning Signs of Acute Risk:

- Threatening to hurt or kill him or herself, or talking of wanting to hurt or kill him/herself; and/or,
- Looking for ways to kill him/herself by seeking access to firearms, available pills, or other means; and/or,
- Talking or writing about death, dying or suicide, when these actions are out of the ordinary.

These might be remembered as expressed or communicated **IDEATION**. If observed, seek help as soon as possible by contacting a mental health professional or calling 1-800-273-TALK (8255) for a referral.

Additional Warning Signs:

- Increased **SUBSTANCE** (alcohol or drug) use
- No reason for living; no sense of **PURPOSE** in life
- **ANXIETY**, agitation, unable to sleep or sleeping all the time
- Feeling **TRAPPED** - like there's no way out
- **HOPELESSNESS**
- **WITHDRAWING** from friends, family and society
- Rage, uncontrolled **ANGER**, seeking revenge
- Acting **RECKLESS** or engaging in risky activities, seemingly without thinking
- Dramatic **MOOD** changes

If observed, seek help as soon as possible by contacting a mental health professional or calling 1-800-273-TALK (8255) for a referral.

These warning signs were compiled by a task force of expert clinical-researchers and 'translated' for the general public.



**SAFETY PLAN: VA VERSION**

**Step 1: Warning signs:**

1. \_\_\_\_\_
2. \_\_\_\_\_
3. \_\_\_\_\_

**Step 2: Internal coping strategies - Things I can do to take my mind off my problems without contacting another person:**

1. \_\_\_\_\_
2. \_\_\_\_\_
3. \_\_\_\_\_

**Step 3: People and social settings that provide distraction:**

1. Name \_\_\_\_\_ Phone \_\_\_\_\_
2. Name \_\_\_\_\_ Phone \_\_\_\_\_
3. Place \_\_\_\_\_ 4. Place \_\_\_\_\_

**Step 4: People whom I can ask for help:**

1. Name \_\_\_\_\_ Phone \_\_\_\_\_
2. Name \_\_\_\_\_ Phone \_\_\_\_\_
3. Name \_\_\_\_\_ Phone \_\_\_\_\_

**Step 5: Professionals or agencies I can contact during a crisis:**

1. Clinician Name \_\_\_\_\_ Phone \_\_\_\_\_  
Clinician Pager or Emergency Contact # \_\_\_\_\_
2. Clinician Name \_\_\_\_\_ Phone \_\_\_\_\_  
Clinician Pager or Emergency Contact # \_\_\_\_\_
3. Local Urgent Care Services \_\_\_\_\_  
Urgent Care Services Address \_\_\_\_\_  
Urgent Care Services Phone \_\_\_\_\_
4. VA Suicide Prevention Resource Coordinator Name \_\_\_\_\_  
VA Suicide Prevention Resource Coordinator Phone \_\_\_\_\_
5. VA Suicide Prevention Hotline Phone: 1-800-273-TALK (8255), push 1 to reach a VA mental health clinician

**Step 6: Making the environment safe:**

1. \_\_\_\_\_
2. \_\_\_\_\_

## VA Safety Plan: Brief Instructions\*

### Step 1: Recognizing Warning Signs

- Ask “How will you know when the safety plan should be used?”
- Ask, “What do you experience when you start to think about suicide or feel extremely distressed?”
- List warning signs (thoughts, images, thinking processes, mood, and/or behaviors) using the patients’ own words.

### Step 2: Using Internal Coping Strategies

- Ask “What can you do, on your own, if you become suicidal again, to help yourself not to act on your thoughts or urges?”
- Ask “How likely do you think you would be able to do this step during a time of crisis?”
- If doubt about using coping strategies is expressed, ask “What might stand in the way of you thinking of these activities or doing them if you think of them?”
- Use a collaborative, problem solving approach to ensure that potential roadblocks are addressed and/or that alternative coping strategies are identified.

### Step 3: Social Contacts Who May Distract from the Crisis

- Instruct patients to use Step 3 if Step 2 does not resolve the crisis or lower risk.
- Ask “Who or what social settings help you take your mind off your problems at least for a little while? “Who helps you feel better when you socialize with them?”
- Ask patients to list several people and social settings, in case the first option is unavailable.
- Ask for safe places they can go to do be around people, e.g. coffee shop.
- Remember, in this step, suicidal thoughts and feelings are not revealed.

### Step 4: Contacting Family Members or Friends Who May Offer Help to Resolve a Crisis

- Instruct patients to use Step 4 if Step 3 does not resolve the crisis or lower risk.
- Ask “Among your family or friends, who do you think you could contact for help during a crisis?” or “Who is supportive of you and who do you feel that you can talk with when you’re under stress?”
- Ask patients to list several people, in case they cannot reach the first person on the list. Prioritize the list. In this step, unlike the previous step, patients reveal they are in crisis.
- Ask “How likely would you be willing to contact these individuals?”
- If doubt is expressed about contacting individuals, identify potential obstacles and problem solve ways to overcome them.

### Step 5: Contacting Professionals and Agencies

- Instruct patients to use Step 5 if Step 4 does not resolve the crisis or lower risk.
- Ask “Who are the mental health professionals that we should identify to be on your safety plan?” and “Are there other health care providers?”
- List names, numbers and/or locations of clinicians, local urgent care services, VA Suicide Prevention Coordinator, VA Suicide Prevention Hotline (1-800-273-TALK (8255))
- If doubt is expressed about contacting individuals, identify potential obstacles and problem solve ways to overcome them.

### Step 6: Reducing the Potential for Use of Lethal Means

- The clinician should ask patients which means they would consider using during a suicidal crisis and collaboratively identify ways to secure or limit access to these means.
- For methods with low lethality, clinicians may ask veterans to remove or restrict their access to these methods themselves.
- Restricting the veterans’ access to a highly lethal method should be done by a designated, responsible person—usually a family member or close friend, or the police.

\*See Safety Plan Treatment Manual to Reduce Suicide Risk: Veteran Version (Stanley & Brown, 2008) for a full description of the instructions.



## The Unique Challenges of Military Families

Center for Deployment Psychology  
Uniformed Services University of the Health Sciences



## Disclaimer

The views expressed are those of the presenters and do not necessarily reflect the opinions of the Uniformed Services University of the Health Sciences, the Department of Defense, or the U.S. Government.



## Learning Objectives

1. Discuss factors that impact marriage, divorce, and relationships in military families.
2. Recognize stressors and risk factors for military families throughout the stages of the deployment cycle.
3. Recognize mechanisms of resilience and protective factors for military families throughout the stages of the deployment cycle.
4. Identify relevant themes for therapy with military families through the stages of the deployment cycle.





## Agenda

- The Definition of the Military Family
- Military Marriages
- Military Family Life
- The Impact of the Deployment Cycle on Military Families
- Resources




## The Definition of the Military Family

*“When one person joins the military  
the whole family serves.”*



## Family Demographics

### Total Force Military Personnel & Family Members



## Military Families

- Dependents
  - Spouses \*\*
  - Children
  - Stepchildren
- Non-Dependents
  - Parents \*
  - Siblings \*
  - Extended family
  - Unmarried partners
  - Adult children





## Military Marriages

*"If the Army wanted you to have a wife, they would have issued you one."*




9



## Marital Status

Total Force Marital Status



2013 Demographics Profile of the Military Community (2014)



10



## Military Marriages

Unique attributes of military marriages:

- When the military calls, the Service member must go
- Separation is standard
- Reunification is also a normal occurrence
- Separation from family/friends is common

Devries et al. (2012)



11



## Marriage and Divorce in the Military



Photo by AzureCitizen, image courtesy of: [http://en.wikipedia.org/wiki/GNU\\_Free\\_Documentation\\_License](http://en.wikipedia.org/wiki/GNU_Free_Documentation_License)

- Young marriages
- Marriage/divorce rates
- Impact of deployment
- Marital quality
- Infidelity

Adler-Baeder et al. (2005) Hogan & Seifert (2010); Karney & Crown (2007); Karney et al. (2012); Negrusa et al. (2014); Riviere et al. (2012); Schumm et al. (2012); Snyder et al. (2012)



12



## Increased Hazard of Divorce

- Greater cumulative time deployed
- Couples married before 9/11
- Effect of deployment greater for:
  - Female Service members
  - Service members in dual-military marriages



Negrusa et al. (2014)



13



## Marital Quality

Examined 2003-2009 trends in marital functioning indicators:

- Marital quality declined
- Reports of past-year infidelity increased
- Reports of separation/divorce intent increased
- *No increases observed* in marital dissolution rates




Riviere et al. (2012)



14



## Impact of Deployment on Couples



Milliken et al. (2007)



15



## Evidence-Based Couples Counseling in the DoD/VA

- Gottman Method Couples Therapy
  - <http://www.gottman.com>
- Emotionally Focused Therapy for Couples (EFT)
  - <http://www.iceeft.com>
- Integrative Behavioral Couples Therapy (IBCT)
  - <http://ibct.psych.ucla.edu>
- Cognitive-Behavioral Conjoint Therapy for PTSD (CBCT) \*
  - <http://www.coupletherapyforptsd.com>



16



## Military Families

*“Military families are families with unique challenges.”*




17



## Parental Status

### Total Force Parental Status



2013 Demographics Profile of the Military Community (2014)



18



## “Greedy Institutions”



Both the military and families demand:

- Commitment
- Loyalty
- Time
- Energy

Blaisure et al. (2012); Segal (1986)



19



## Family Stressors

### Normative

- Occur for most families
- Expected

### Normative Military

- Occur for most **military** families
- Expected

### Catastrophic

- Do not occur to most families
- Unexpected

Blaisure et al. (2012)



20



## Normative Stressors of Military Family Life

- Frequent relocations (PCS)
- Spouse employment opportunities
- Separations
- Deployments
- Risk of injury or death



Image courtesy of: <http://www.dodlive.mil/index.php/2011/06/how-to-ease-the-summertime-move/>

Blaisure et al. (2012); Lim & Schulker (2010)



21



*If you knew you had to move your entire household every 2-3 years, how would you live your life differently than you do now?*

Blaisure et al. (2012), p. 57



22



## 2014 Blue Star Families Survey

### Top Stressors Associated with the Military

- 69% = Deployments
- 51% = Isolation from family & friends

### Military Spouse Employment

- 84% (employed) = military lifestyle negatively impacted pursuit of career
- 53% (unemployed) = would like to be employed outside the home

Blue Star Families (2014)



23



## Normative Military Stressors for Children

- Relocation
- Education
- Child Care
- Deployments



Blaisure et al. (2012)



24



## Resiliency in Military Children

- Sense of belonging/community
- Adaptable
- Tolerant of diversity
- Responsible/independent
- Respect for authority



Easterbrooks et al. (2013); Hall (2012); Park (2011)




25



## Blended Families

- Children of different developmental ages and stages
- Higher risk of emotional / behavioral problems
- Torn between biological parents
- Two households
- Issues of isolation



Adler-Baeder et al. (2005); Hall (2008)



27



## LGB Families

- 20 September 2011 = Repeal of DADT
- 26 June 2013 = Repeal of Section 3 of DOMA



- Prejudice/ Discrimination
- Identity Concealment
- Relocation Issues
- Social Support

Ender et al. (2012); Herek (2009); Oswald & Sternberg (2014)



28



## Keys to Family Resilience

### Family Belief Systems

Making meaning of adversity

Positive outlook

Transcendence & spirituality

### Family Organizational Patterns

Flexibility

Connectedness

Social & economic resources

### Family Communication Processes

Clarity

Open emotional expression

Collaborative problem solving

Walsh (2006)



29



## Mechanisms of Resilience in Military Families

- Psychoeducation & developmental guidance
- Shared family narratives
- Open & effective communication
- Family resiliency (coping) skills
- Effective & coordinated parent leadership

Saltzman et al. (2014)



31



## Resiliency Training for Military Families

- FOCUS (Families OverComing Under Stress)
  - <http://www.focusproject.org>
  - <http://www.focusworld.org>
- HomeFront Strong
  - <http://m-span.org/programs-for-military-families/homefront-strong>



32




## The Deployment Cycle & Its Impact on Family Members



35



## Emotional Cycle of Deployment



Pincus et al. (2001)



36

**CDP**

## MilSpouse Cycle of Deployment

shoot he's leaving again  
rumor  
mill

Pre-Deployment

field training #/5

Post-Deployment

MilSpouse

field training #/10

Deployment

emotionally unstable

denial

field training #/2  
field training #/1  
get used to living together again

Deployment Cycle

Redeployment

ooooo la la!  
excited!

Sustainment

ck, i got this  
bitchy me bitchkins

Image courtesy of: Patricia Santiago (2010)

hjf 37

**CDP**

## Case Study: Alison Talks about Deployments

hjf 38

**CDP**

## Parenting Cycle of Deployment

Pre-Deployment	Deployment	Post-Deployment
Looking Ahead	Parenting (War Zone / Home Front)	Facing Reality
Saying Goodbye	Surviving the Home Stretch	Moving Forward

DeVoe & Ross (2012)

hjf 39

**CDP**

## Pre-Deployment Stage

*"A holding pattern during which life cannot yet begin."*


DeVoe & Ross (2012)

hjf 40





## Emotional Reactions



DeVoe & Ross (2012)



41



## Parenting

- Looking Ahead
  - How/when to communicate with children about impending departure
  - Handling feelings & responding to children
- Saying Goodbye
  - Importance of saying goodbye face-to-face

Pre-Deployment

DeVoe & Ross (2012)



42



## Helping Families

- Facilitate communication
  - Sharing of feelings
  - Discussion of expectations
- Foster connection
- Teach positive coping skills
- Emphasize the importance of support
- Help couples address practical issues
- Discuss parenting issues

Pre-Deployment

Pavlicin (2003); Pincus et al. (2001)



43



## Helping Children

- Encourage quality time with each child
- Emphasize honesty
- Help parents provide reassurance
- Foster connection
- Create communication plan
- Facilitate conversations with schools/teachers



Dod photo by PO2 Stephanie Tigner/Released

Pre-Deployment

Military OneSource (2013); Pavlicin (2003)



44



## Themes for Therapy

- Affective Education
- Exploring Roles
- Exploring Unresolved Conflicts
- Expectations of Fidelity



Pre-Deployment



## Deployment Stage

*“Repeat after me ... I can do this.”*



## Emotional Reactions

- Deployment
  - Disorientation, mixed emotions
  - Relief that anticipation is over
- Sustainment
  - Self-growth, independence
  - Less angry, but lonelier
- Re-Deployment (Anticipation of Return)
  - Excitement, apprehension

Deployment



## Parenting

- Parenting from the Home Front
  - Increased responsibilities = increased stress
- Parenting from a War Zone
  - Relinquishment of daily involvement
  - Adaptation of new strategies
- Surviving the Home Stretch
  - How will children respond to reunion?

Deployment



## Deployment Protective Factors

- Family readiness
- Active coping styles
- “Making meaning” of the deployment
- Strong community of social support
- Acceptance of military lifestyle
- Optimism
- Self-reliance
- Ability to adopt flexible gender roles

Hammer et al. (2006); Patterson & McCubbin (1984); Rosen et al. (1993); Walsh (2006); Weins & Boss (2006)



49



## Family Risk Factors During Deployment

- Rigid coping style
- History of family dysfunction/ behavioral health problems
- Families experiencing first military separation
- New to duty location
- Young families
- Pregnant spouses
- Single parent families
- Dual military families
- Families with foreign spouses
- Blended families
- Selected Reserve

Blount et al. (1992); Darwin (2012); Hall (2008) Huffman & Payne (2006); Kelley (2006); Weins & Boss (2006); Wolpert et al. (2000)



50



## Impact of Deployment on Children

- Emotional / Behavioral Difficulties
- Cumulative Months of Deployment
- Child Maltreatment



US Army photo by PFC Jessica M. Kuhn. Image courtesy of <https://creativecommons.org/licenses/by/3.0/legalcode>

Deployment

Chandra et al. (2011); Flake et al. (2009); Gibbs et al. (2007); Lester et al. (2010); Rentz et al. (2007)



51



## Parental Deployment & Adolescent Mental Health

Reporting of any familial deployment (parent or sibling) was associated with increased odds of experiencing:

- Sadness/hopelessness
- Depressive symptoms
- Suicidal ideation



Cederbaum et al. (2013)



52



## 2014 Blue Star Families Survey

### Top 5 **Positive** Impacts of Deployment on Children

- Adaptability
- Increased Independence
- Personal Growth
- Increased Resilience
- Increased Pride

### Top 5 **Negative** Impacts of Deployment on Children

- Separation Anxiety
- Worry
- Irritability
- Difficulty Sleeping
- Difficulty Concentrating



## Case Study: Kayla Talks about Deployments



## Helping Families

- Facilitate setting personal goals
- Encourage creation of a support network
- Teach stress management techniques
- Highlight the importance of self-care



Deployment



## Helping Children

- Encourage consistency & routine
- Highlight the importance of support from other adults
- Help parents model self-care
- Facilitate good communication
- Emphasize the importance of fun
- Remind parents to let kids be kids

Deployment



**CDP** **Case Study: Kayla's Experience During Deployment**



hjf  58

**CDP** **Themes for Therapy**

- Deployment:  
Stress Reduction
- Sustainment:  
Resilience & Growth
- Re-Deployment:  
Expectation Management



Deployment

Laser & Stephens (2012) hjf  59

**CDP**

**Post-Deployment Stage**

*"Hello, Stranger.  
Let's cohabit and raise these kids."*

Sanderlin (2012) hjf  60


**CDP** **Emotional Reactions**

Reunion → Reintegration

- Elation
- Relief
- Frustration


- Tension
- Resentment
- Comfort

Post-Deployment

Logan (1987); Pincus, et al. (2001) hjf  61

**CDP**

## Case Study: Homecoming



62

**CDP**

## Parenting

- Facing Reality
  - Reconnect & rebuild parent-child relationships
  - Role negotiation / co-parenting challenges
- Moving Forward
  - Re-establish family equilibrium
  - Develop new parenting routines
  - Incorporate legacy of deployment

Post- Deployment


DeVoe & Ross (2012)

hjf **US** 63

**CDP**

## The Impact of Mental Health Issues

- Depression, PTSD
- Caregiver Burden
- “As goes the parent, so goes the child”




Blaisure et al. (2012); Blow et al. (2013); Flake et al. (2009); Goff et al. (2007); Lester et al. (2010); Saltzman, et al. (2014); Savers et al. (2009); Tanielian et al. (2013)

hjf **US** 65

**CDP**

## Case Study: Alison Talks about James' Return Home



66

**CDP** **Case Study: Kayla Talks about Her Dad's Return Home**



67

**CDP** **Facilitating Successful Reunions**

Service Member's Expectations	Partner's Expectations	Therapist's Role (how to help)
Isolate vs. great sex 24/7	Fairy-tale reunion	<b>Set realistic expectations</b>
Immediate return to "old ways"	New routines	<b>Remind that adjustment takes time</b>
Separated from family	Earned a break	<b>Stress that both made sacrifices</b>
No one understands	Drill with questions	<b>Teach communication skills</b>

Military OneSource (2012); Pavlicin (2003)

hjf **CDP** 68

**CDP** **Successful Reunions with Children**

Things to remember:

- Children react differently to homecoming depending on their age & relationship with deployed parent
- Children are often loyal to the parent that stays behind
- Children may feel anxious about the Service member leaving again
- Children will wonder if rules at home will change

Post- Deployment

hjf **CDP** 69

Pavlicin (2003); Pincus et al. (2001)

**CDP** **Child Development and Reunion & Reintegration**

Children respond differently to homecomings and parental reintegration depending on their age and developmental stage.

**REUNION & REINTEGRATION WITH CHILDREN**

**Infants/Toddlers (Ages 0-3)**  
 Infancy is a time of rapid growth and development (both physically and cognitively). Attachment is the major developmental milestone during this stage.

**Homecoming Reactions**

- "Stranger Reaction"
- Increased sensitivity
- Crying, clinginess, disrupted schedule
- Delayed milestones
- Temper tantrums
- Nightmares
- Regression in skills

**Facilitating Reintegration**

- Realistic expectations
- Direct contact
- Consistency
- Opportunities for connection
- Take things slowly
- Expect regression

Post- Deployment

hjf **CDP** 70



## Themes for Therapy

- Revisiting Family Roles
- Opening Communication
- Creating Opportunities for Appreciation and Caring



Post- Deployment

Laser & Stephens (2012); image by: Scott Maxwell, <http://thegoldguys.blogspot.com>.  
Image courtesy of: <https://creativecommons.org/licenses/by-sa/2.0/legalcode>



72



## Online Resources

Please see  
**Military Families  
Online Resources  
Handout**  
for additional military  
family resources.



79



## CDP Website: Deploymentpsych.org

### Features include:

- Descriptions and schedules of upcoming training events
- Blog updated daily with a range of relevant content
- Articles by subject matter experts related to deployment psychology, including PTSD, mTBI, depression, and insomnia
- Other resources and information for behavioral health providers
- Links to CDP's Facebook page and Twitter feed



80



## Online Learning

The following online courses are located on the CDP website at:

<http://www.deploymentpsych.org/content/online-courses>

**NOTE: All of these courses can be taken for free or for CE Credits for a fee**


- Cognitive Processing Therapy (CPT) for PTSD in Veterans and Military Personnel (1.25 CE Credits)
- Prolonged Exposure Therapy for PTSD in Veterans and Military Personnel (1.25 CE Credits)
- Epidemiology of PTSD in Veterans: Working with Service Members and Veterans with PTSD (1.5 CE Credits)
- Provider Resiliency and Self-Care: An Ethical Issue (1 CE Credit)
- Military Cultural Competence (1.25 CE Credits)
- The Impact of Deployment and Combat Stress on Families and Children, Part 1 (2.25 CE Credits)
- The Impact of Deployment and Combat Stress on Families and Children, Part 2 (1.75 CE Credits)
- The Fundamentals of Traumatic Brain Injury (TBI) (1.5 CE Credits)
- Identification, Prevention, & Treatment of Suicidal Behavior in Service Members & Veterans (2.25 CE Credits)
- Depression in Service Members and Veterans (1.25 CE Credits)

*All of these courses and several others are contained in the Serving Our Veterans Behavioral Health Certificate program, which also includes 20+ hours of Continuing Education Credits for \$350.*



81







## Provider Support

CDP's "Provider Portal" is exclusively for individuals trained by the CDP in evidence-based psychotherapies (e.g., CPT, PE, and CBT-I)


**Features include:**

- Consultation message boards
- Hosted consultation calls
- Printable fact sheets, manuals, handouts, and other materials
- FAQs and one-on-one interaction with answers from SMEs
- Videos, webinars, and other multimedia training aids

Participants in CDP's evidence-based training will automatically receive an email instructing them how to activate their user name and access the "Provider Portal" section at Deploymentpsych.org.


82




## How to Contact Us

**Center for Deployment Psychology**  
 Department of Medical & Clinical Psychology  
 Uniformed Services University of the Health Sciences  
 4301 Jones Bridge Road, Executive Office: Bldg. 11300-602  
 Bethesda, MD 20813-4768


**Email:** [General@DeploymentPsych.org](mailto:General@DeploymentPsych.org)  
**Website:** [DeploymentPsych.org](http://DeploymentPsych.org)  
**Facebook:** <http://www.facebook.com/DeploymentPsych>  
**Twitter:** @DeploymentPsych




83



## Resources




73




## Military OneSource

- Confidential services available via telephone & online
- Comprehensive information on every aspect of military life:
  - Deployment
  - Reunion
  - Relationships
  - Grief
  - Spouse employment & education
  - Parenting & childhood



<http://www.militaryonesource.mil>



74



## FOCUS: Families OverComing Under Stress

- Training in core resilience skills
  - Emotion regulation
  - Communication
  - Problem solving
  - Goal-setting
  - Managing deployment reminders



Image courtesy of: <http://www.dodlive.mil/index.php/2010/12/medical-monday-project-focus/>

- FOCUS World (online resiliency training)
- FOCUS On the Go! (mobile app)

<http://www.focusproject.org>



33



## FOCUS World



FOCUS World ([www.focusworld.org](http://www.focusworld.org)) is an interactive, online educational tool that helps military families become stronger in the face of challenges by providing both parents and kids a fun place to learn and practice important skills, such as:

<http://www.focusworld.org>



34



## Military Kids Connect

- Online community for military children (ages 6-17)
- Access to age-appropriate resources to support children dealing with the challenges of military life
- Additional resources for parents & teachers



<http://www.militarykidsconnect.dcoe.mil>



75



## Military Kids Connect



<http://www.militarykidsconnect.dcoe.mil>



76



## Sesame Street Toolkit: Talk, Listen, Connect


- DVDs: Deployment, Homecomings, Grieving
- Website: Military Families Near & Far
- Mobile Apps:
  - Sesame Street for Military Families
  - The Big Moving Adventure



<http://www.sesamestreet.org/parents/topicsandactivities/toolkits/tlc>



77



## Department of Defense 2013 Demographics Profile of the Military Community

<http://www.militaryonesource.mil>  
Click on "Reports and Surveys"



78

## The Unique Challenges of Military Families

### References

- Adler-Baeder, F., Pittman, J. F., & Taylor, L. (2005). The prevalence of marital transitions in military families. *Journal of Divorce & Remarriage*, 44(1-2), 91-106.  
[http://dx.doi.org/10.1300/J087v44n01\\_05](http://dx.doi.org/10.1300/J087v44n01_05)
- Alvarez, L. (2006, July 8). An internet lifeline for troops in Iraq and loved ones at home. *The New York Times*. Retrieved from:  
[http://www.nytimes.com/2006/07/08/us/08FAMILY.html?pagewanted=1&\\_r=2](http://www.nytimes.com/2006/07/08/us/08FAMILY.html?pagewanted=1&_r=2)
- Barnett, J. E., Getkin, C., Grover, A., King, D., Schroedle, A., Sherman, M. D., & Tews, R. (2012). Children's reactions to deployment, reintegration, and injuries by age groups, with recommendations on how to help them. In N. D. Ainspan & W. E. Penk (Eds.), *When the warrior returns: Making the transition at home* (pp. 165-184). Annapolis, MD: Naval Institute Press.
- Blaisure, K. R., Saatoff-Wells, T., Pereira, A., Wadsworth, S. M., & Dombro, A. L. (2012). *Serving military families in the 21<sup>st</sup> century*. New York, NY: Routledge.
- Blount, B. W., Curry, A., & Lubin, G. I. (1992). Family separations in the military. *Military Medicine*, 157(2), 76-80.
- Blow, A. J., Gorman, L., Ganoczy, D., Kees, M., Kashy, D. A., Valenstein, M., ... Chermack, S. (2013). Hazardous drinking and family functioning in National Guard veterans and spouses postdeployment. *Journal of Family Psychology*, 27(2), 303-313.  
<http://dx.doi.org/10.1037/a0031881>
- Blue Star Families. (2014). *2014 military families lifestyle survey: Comprehensive report*. Retrieved from:  
[http://bluestarfam.org/sites/default/files/media/stuff/bsf\\_report\\_comprehensive\\_reportfinal\\_single\\_pages.pdf](http://bluestarfam.org/sites/default/files/media/stuff/bsf_report_comprehensive_reportfinal_single_pages.pdf)
- Bowles, S. V., Pollock, L. D., Cato, C., Moore, M., Wadsworth, S. M., Anagnostopoulos, V., ... Bates, M. J. (2012). Building resilience in the military family during and following deployment. In N. D. Ainspan & W. E. Penk (Eds.), *When the warrior returns: Making the transition at home* (pp. 79-99). Annapolis, MD: Naval Institute Press.
- Cederbaum, J.A., Gilreath, T.D., Benbenishty, R. Astor, R.A., Pineda, D., DePedro, K.T., ... Stuel, H. (2014). Well-being and suicidal ideation of secondary school students from military families. *Journal of Adolescent Health*, 54(6), 672-677. <http://dx.doi.org/10.1016/j.jadohealth.2013.09.006>
- Chandra, A., Lara-Cinisomo, S., Jaycox, L. H., Tanielian, T., Han, B., Burns, R. M., & Ruder, T. (2011). *Views from the homefront: The experiences of youth and spouses from military families* (RAND Document No. TR-913-NMFA). Santa Monica, CA: The RAND Corporation.
- Darwin, J. (2012). Families of National Guard and Reserve service members. In N. D. Ainspan & W. E. Penk (Eds.), *When the warrior returns: Making the transition at home* (pp. 242-252). Annapolis, MD: Naval Institute Press.
- Derenne, J. L. (2008). *Raising an emotionally healthy child: How not to be a helicopter parent* [PowerPoint slides]. Retrieved from  
<http://www.mcw.edu/FileLibrary/User/llemahie/FacultyAffairs/WomensFacultyCouncil/Derenn e.RaisinganEmotionallyHe.ppt>

## The Unique Challenges of Military Families

- DeVoe, E. R. & Ross, A. (2012). The parenting cycle of deployment. *Military Medicine*, 177(2), 184-190.
- Devries, M. R., Hughes, H. K., Watson, H, & Moore, B. A. (2012). Understanding the military culture. In B. A. Moore (Ed.), *Handbook of counseling military couples* (pp. 7-18). New York, NY: Routledge.
- Dimidjian, S., Martell, C. R., & Christensen, A. (2008). Integrative behavioral couple therapy. In A. S. Gurman (Ed.), *Clinical handbook of couple therapy* (4<sup>th</sup> ed., pp. 73-103). New York, NY: The Guilford Press.
- Easterbrooks, M.A., Ginsburg, K., & Lerner, R.M. (2013). Resilience among military youth. *The Future of Children*, 23(2), 99-120.
- Ender, M.G., Rohall, D. E., Brennan, A. J., Matthews, M. D., & Smith, I. (2012). Civilian, ROTC, and military academy undergraduate attitudes toward homosexuals in the U.S. military: A research note. *Armed Forces & Society*, 38(1), 164-172. <http://dx.doi.org/10.1177/0095327X11410856>
- Flake, E. M., Davis, B. E., Johnson, P. L., & Middleton, L. S. (2009). The psychosocial effects of deployment on military children. *Journal of Developmental Behavioral Pediatrics*, 30(4), 271-278. <http://dx.doi.org/10.1097/DBP.0b013e3181aac6e4>
- Gibbs, D. A., Martin, S. L., Kupper, L. L., & Johnson, R. E. (2007). Child maltreatment in enlisted soldiers' families during combat-related deployments. *Journal of the American Medical Association*, 298(5), 528-535. <http://dx.doi.org/10.1001/jama.298.5.528>
- Goff, B. S. N., Crow, J. R., Reisberg, A. M. J., & Hamilton, S. (2007). The impact of individual trauma symptoms of deployed soldiers on relationship satisfaction. *Journal of Family Psychology*, 21(3), 344-353. <http://dx.doi.org/10.1037/0893-3200.21.3.344>
- Gottman, J. M. & Gottman, J. S. (2008). Gottman method couple therapy. In A. S. Gurman (Ed.), *Clinical handbook of couple therapy* (4<sup>th</sup> ed., pp. 138-164). New York, NY: The Guilford Press.
- Gottman, J. M. & Silver, N. (1999). *The seven principles for making marriage work*. New York, NY: Three Rivers Press.
- Hall, L. K. (2008). *Counseling military families: What mental health professionals need to know*. New York, NY: Routledge.
- Hall, L.K. (2012). The military lifestyle and the relationship. In B.A. Moore (Ed.), *Handbook of counseling military couples* (pp. 137-156). New York, NY: Routledge.
- Hammer, L. B., Cullen, J. C., Marchand, G. C., & Dezsofi, J. A. (2006). Reducing the negative impact of work-family conflict on military personnel: Individual coping strategies and multilevel interventions. In C. A. Castro, A. B. Adler, & T. W. Britt (Eds.), *Military life: The psychology of serving in peace and combat, Volume 3: The military family* (pp. 220-242). Bridgeport, CT: Praeger Security International.
- Herek, G. M. (2009). Hate crimes and stigma-related experiences among sexual minority adults in the United States: Prevalence estimates from a national probability sample. *Journal of Interpersonal Violence*, 24(1), 54-74. <http://dx.doi.org/10.1177/0886260508316477>

## The Unique Challenges of Military Families

- Hogan, P. F. & Seifert, R. F. (2010). Marriage and the military: Evidence that those who serve marry earlier and divorce earlier. *Armed Forces & Society*, 36(3), 420-438.  
<http://dx.doi.org/10.1177/009537X09351228>
- Huffman, A. H. & Payne, S. C. (2006). The challenges and benefits of dual-military marriages. In C. A. Castro, A. B. Adler, & T. W. Britt (Eds.), *Military Life: The psychology of serving in peace and combat, Volume 3: The military family* (pp. 115-137). Bridgeport, CT: Praeger Security International.
- Johnson, S. M. (2008). Emotionally focused couple therapy. In A. S. Gurman (Ed.), *Clinical handbook of couple therapy* (4<sup>th</sup> ed., pp. 107-137). New York, NY: The Guildford Press.
- Karney, B. R. & Crown, J. S. (2007). *Families under stress: An assessment of data, theory, and research on marriage and divorce in the military* (RAND Document No. MG-599-OSD). Santa Monica, CA: The RAND Corporation.
- Karney, B. R., Loughran, D. S., & Pollard, M. S. (2012). Comparing marital status and divorce status in civilian and military populations. *Journal of Family Issues*, 33(12), 1572-1594.  
<http://dx.doi.org/10.1177/0192513X12439690>
- Kelley, M. L. (2006). Single military parents in the new millennium. In C. A. Castro, A. B. Adler, & T. W. Britt (Eds.), *Military life: The psychology of serving in peace and combat, Volume 3: The military family* (pp. 93-114). Bridgeport, CT: Praeger Security International.
- Laser, J. A. & Stephens, P. M. (2011). Working with military families through deployment and beyond. *Clinical Social Work Journal*, 39, 28-38. <http://dx.doi.org/10.1007/s10615-010-0310-5>
- Lester, P., Peterson, K., Reeves, J., Knauss, L., Glover, D., Mogil, C., ... Beardslee, W. (2010). The long war and parental combat deployment: Effects on military children and at-home spouses. *Journal of the American Academy of Child & Adolescent Psychiatry*, 49(4), 310-320.
- Lester, P., Saltzman, W.R., Woodward, K., Glover, D., Leskin, G.A., Bursch, B., ... Beardslee, W. (2012). Evaluation of a family-centered prevention intervention for military children and families facing wartime deployments. *American Journal of Public Health*, 102, S48-S54.  
<http://dx.doi.org/10.2015/AJPH.2010.30088>
- Lim, N & Schulker, D. (2010). *Measuring underemployment among military spouses* (RAND Document No. MG-918-OSD). Santa Monica, CA: The RAND Corporation.
- Logan, K. V. (1987). The emotional cycle of deployment. *U.S. Naval Institute Proceedings*, 113, 43-47.
- Military OneSource. (2013). *Military deployment guide: Preparing you and your family for the road ahead*. Retrieved from  
<http://www.militaryonesource.mil/12038/MOS/ResourceGides/DeploymentGuide.pdf>
- Milliken, C.S., Auchterlonie, J.L., & Hoge, C.W. (2007). Longitudinal assessment of mental health problems among active and reserve component soldiers returning from the Iraq war. *Journal of the American Medical Association*, 298(18), 2141-2148.  
<http://dx.doi.org/10.1001/jama.298.18.2141>

## The Unique Challenges of Military Families

- Monson, C.M., Fredman, S.J., & Adair, K.C. (2008). Cognitive-behavioral conjoint therapy for PTSD: Application to Operation Enduring and Iraqi Freedom veterans. *Journal of Clinical Psychology: In Session, 64*(8), 958-971.
- Morris, J. (2008, October 23). *MN ARNG reintegration initiative: Warrior transition: From warrior-citizen to citizen-soldier* [PowerPoint slides]. Lecture presented at *Topics for Deployment Psychology*. Center for Deployment Psychology, Bethesda, MD.
- Negrusa, S., Negrusa, B., & Hosek, J. (2014). Gone to war: Have deployments increased divorces? *Journal of Population Economics, 27*(2), 473-496. <http://dx.doi.org/10.1007/S00148-013-0485-5>
- Office of the Deputy Assistant Secretary of Defense, Military Community & Family Policy. (2014). *2013 demographics profile of the military community*. Retrieved from: <http://www.militaryonesource.mil/12038/MOS/Reports/2013-Demographics-Report.pdf>
- Oswald, R.F., & Sternberg, M. M. (2014). Lesbian, gay, and bisexual military families: Visible but legally marginalized. In S.M. Wadsworth & D. S. Riggs (Eds.), *Military deployment and its consequences for families* (pp. 133-147). New York, NY:Springer.
- Park, N. (2011). Military children and families: Strengths and challenges during peace and war. *American Psychologist, 66*(1), 65-72. <http://dx.doi.org/10.1037/a0021249>
- Patterson, J. M & McCubbin, H. I. (1984). Gender roles and coping. *Journal of Marriage & the Family, 46*(1), 95-104. <http://dx.doi.org/10.2307/351868>
- Pavlicin, K. M. (2003). *Surviving deployment: A guide for military families*. St. Paul, MN: Elsa Reva Publishing.
- Pincus, S. H., House, R., Christenson, J., & Adler, L. E. (2001). The emotional cycle of deployment: A military family perspective. *U.S. Army Medical Department Journal, Apr-Jun, 15-23*.
- Rentz, E. D., Marshall, S. W., Loomis, D., Casteel, C., Martin, S. L., & Gibbs, D.A. (2007). Effect of deployment on the occurrence of child maltreatment in military and nonmilitary families. *American Journal of Epidemiology, 165*(10), 1199-1206. <http://dx.doi.org/10.1093/aje/kwm008>
- Riviere, L.A., Merrill, J.C., Thomas, J.L., Wilk, J.E., & Bliese, P.D. (2012). 2003-2009 marital functioning trends among U.S. enlisted soldiers following combat deployments. *Military Medicine, 177*(10), 1169-1177.
- Rosen, L.N., Teitelbaum, J.M., & Westhuis, D.J. (1993). Stressors, stress mediators, and emotional well-being among spouses of soldiers deployed to the Persian Gulf during Operation Desert Shield/Storm. *Journal of Applied Social Psychology, 23*(19), 1587-1593. <http://dx.doi.org/10.1111/j.1559-1816.1993.tb01049.x>
- Sanderlin, R. (2012, November 13). Finding home again after deployment. *The New York Times*. Retrieved from: <http://atwar.blogs.nytimes.com/2012/11/13/finding-home-again-after-deployment/>
- Saltzman, W.R., Bartoletti, M., Lester, P. & Beardslee, W. (2014). Building resilience in military families. In S. Cozza, M.N. Goldenberg & R.J. Ursano (Eds.), *Care of Military Service Members, Veterans, & their Families*. Arlington, VA: American Psychiatric Association.

## The Unique Challenges of Military Families

- Sayers, S. L., Farrow, V. A., Ross, J., & Oslin, D. W. (2009). Family problems among recently returned military veterans referred for a mental health evaluation. *Journal of Clinical Psychology, 70*(2), 163-170.
- Schumm, W.R., Roy, R.R.N., & Theodore, V. (2012). Separation and divorce. In B.A. Moore (Ed.), *Handbook of counseling military couples* (pp. 157-181). New York, NY: Routledge.
- Segal, M. W. (1986). The military and the family as greedy institutions. *Armed Forces & Society, 13*(1), 9-38. <http://dx.doi.org/10.1177/0095327X8601300101>
- Snyder, D.K., Balderrama-Durbin, C., Fissette, C, Scheider, D.M., Barnett, J.K, & Fiala, S. (2012). Infidelity. In B.A. Moore (Ed.), *Handbook of counseling military couples* (pp. 219-235). New York, NY: Routledge.
- Stanley, T. (2013). The art and science of Gottman method couples therapy. *The Therapist, 25*(2), 18-23.
- Tanielian, T., Ramchand, R. Fisher, M. P., Sims, C. S., Harris, R., & Harrell, M. C. (2013). *Military caregivers: Cornerstones of support for our nation's wounded, ill, and injured veterans* (RAND Document No. RR-244-TEDF). Santa Monica, CA: The RAND Corporation.
- Walsh, F. (2006). *Strengthening family resilience (2<sup>nd</sup> ed)*. New York, NY: The Guildford Press.
- Walter Reed Army Institute of Research. (2011). *Post-deployment resilience training for spouses/couples*.
- Weins, T. W & Boss, P. (2006). Maintaining family resiliency before, during, and after military separation. In C. A. Castro, A. B. Adler, & T. W. Britt (Eds.), *Military life: The psychology of serving in peace and combat: Volume 3: The military family* (pp. 13-38). Bridgeport, CT: Praeger Security International.
- Wolpert, D. S., Martin, J. A., Dougherty, L. M., Rudin, B. J., & Kerner-Hoeg, S. (2000). The special case of the young enlisted family. In J. A. Martin, L. N. Rosen, & L. R. Sparacino (Eds.), *The military family: A practice guide for human service providers* (pp. 43-53). Westport, CT: Praeger.







## Infants/Toddlers (Ages 0-3)

Infancy is a time of rapid growth and development (both physically and cognitively). Attachment is the major developmental milestone during this stage.

### Homecoming Reactions


- “Stranger Reaction”
- Increased sensitivity
- Crying, clinginess, disrupted schedule
- Delayed milestones
- Temper tantrums
- Nightmares
- Regression in skills

### Fostering Reintegration

- Realistic expectations
- Direct contact
- Consistency
- Opportunities for connection
- Take things slowly
- Expect regression

## Pre-schoolers (Ages 3-6)

During the pre-school stage, growth and development continue at a rapid pace. Children perceive that everything revolves around them and magical thinking predominates.



### Homecoming Reactions

- Personalize stress
- Regression in skills
- Acting out
- Time to warm up

### Fostering Reintegration

- Ignore regressive behaviors
- Listen & answer questions
- Opportunities for connection
- Take things slowly



## School Age Children (*Ages 6-12*)

During the school years physical development slows. Children begin to find their place in the world, and they show increases in organization, responsibility, and cognitive and moral development.

### Homecoming Reactions

- Whiny, aggressive, irritable
- Need time to talk
- Need physical attention
- Rapid mood shifts
- Excitement
- Fear

### Fostering Reintegration

- Slowly transition roles/responsibilities
- Include Service member in routines
- One-on-one time
- Connect to school/community resources

---

## Adolescents (*Ages 12-18*)

Adolescence is the time when children prepare for adulthood. Puberty occurs and there is an increased reliance on the peer group to determine norms. Adolescents are now capable of abstract thinking. It is very common for there to be parent/child tension during this time.



### Homecoming Reactions

- Emotionally guarded
- Acting out
- Lower grades
- Relief
- Concern about roles/responsibilities
- Sensitivity

### Fostering Reintegration

- Open communication
- Consistency
- Transition roles/responsibilities
- One-on-one time
- Respect privacy



## **Military Family Online Resources**



### **Military Family Resources:**

Military One Source: <http://www.militaryonesource.mil>

Military Family Research Institute: <https://www.mfri.purdue.edu>

After Deployment: <http://afterdeployment.dcoe.mil>

Real Warriors: <http://www.realwarriors.net>

Yellow Ribbon Program: <http://www.yellowribbon.mil>

Military Partners & Families Coalition: <http://www.milpfc.org>

### **Military Child Resources:**

Military Kids Connect: <https://militarykidsconnect.dcoe.mil>

Talk, Listen, Connect: <http://www.sesamestreet.org/parents/topicsandactivities/toolkits/tlc>

Military Families Near & Far: <https://www.familiesnearandfar.org>

Zero to Three: <http://www.zerotothree.org/>

Military Child Education Coalition: <http://www.militarychild.org/>

Military Youth Coping with Separation Video: <http://bit.ly/1qAiqIO>

### **Evidence-Based Couples Counseling in the DoD/VA:**

Gottman Method Couples Therapy:

<http://www.gottman.com>

<http://csf2.army.mil>

Emotionally Focused Therapy for Couples (EFT):

<http://iceeft.com>

<http://www.strongbonds.org>

Integrative Behavioral Couples therapy (IBCT): <http://www.ibct.psych.ucla.edu>

Cognitive-Behavioral Conjoint Therapy for PTSD (CBCT): <http://www.coupletherapyforptsd.com>

### **Resiliency Training for Military Families:**

Families Overcoming Under Stress (FOCUS): <http://www.focusproject.org/>

FOCUS World: <http://www.focusworld.org>

HomeFront Strong: <http://m-span.org/programs-for-military-families/homefront-strong/>

