Steps to Change PCL Frequency in BHDP

This handout provides detailed instructions and screenshots designed to assist providers in changing the frequency of the PCL in BHDP. This same process can be used to change the frequency of other outcome measures as well.

- 1. Open the patient's BHDP dashboard.
- 2. Click on the wrench icon under the heading "Clinical Section Frequency" as shown below.

🕈 AFP - Search 🥔 Dashboard	× 🗂		
BHOP —	Sparch Dashboard:	Adult BH	Logged in: Ltst, #st[Pt. List User Settings User Guide Help Feedback Reports Logout]
Dashboard Most Recent	Hot Sheet 🚱 All	Clinical Scales	Notes View
PATIENT	TREATMENT HISTORY		😴 ង CARE TEAM MEMBERS
	Treatment Effectiveness Pain Leve		Add Me Name Contact Number S
Add Patient To Patient List :	Clinical Section FREQUENCY	Last Taken Frequency	Click on the
💽 TREATMENT STATUS	Functional Impairment	Every 5 days	Profile Info is not
rimary Diagnosis or Primary Problem	BASIS-24	Every 5 days	Create a Profile
Not Set	PHQ2	Every 60 days	📾 SIGNIFICANT EVENTS +HIP
dditional Diagnosis or Problems	РНОЭ	Every 0 days	No significant events found.
reatment Setting	- GAD2	Every 60 days	
Not Set	GAD7	Every 0 days	C DEPLOYMENT SUMMARY
reatment Modality Not Set	C-SSRS-S Lifetime	Once	Deployment Summary is not available
reatment Type	C-SSRS-S Current	Every 1 days	
Not Set	4Q-PTSD	Every 0 days	GEFLOTMENTS
reatment Frequency Not Set	4QPTSD	Every 60 days	Deployment history is not available
isk for Harm to Self	PCL-C	Every 0 days	MEB INFORMATION -
NotSet	PCL-5	Every 0 days	MEB Information is not available
Not Set	AUDIT-C	Every 60 days	
isk for Being Harmed by Others Not Set (MM/DD/YYY)	AUDIT	Every 0 days	>> Warrior Transition Enrollment
	CSI	Every 365 days	Warrior Transition Enrollment is not available
PATIENT REPORTED WITALS	ISI Screen	Every 5 days	
ain Level:	ISI	Every 0 days	
-SSRS-S: Rehavior? Number:	BAM_Intake	Every 0 days	
Actes Suicide Risk Level	Therapeutic Alliance and Patient Satisfaction	Every 30 days	



- 3. On the "Clinical Section Frequency" screen, scroll down to the PCL.
- 4. Change frequency to "0" to have the patient complete the PCL at every session as shown below. Note that the BHDP varies across services. If you cannot put 0, then put the lowest number possible in.
- 5. Click "Save".



6. Upon return to the "Clinical Section Frequency" on the patient dashboard, verify that the new frequency selection has been saved.

