



National Center for
PTSD
POSTTRAUMATIC STRESS DISORDER

PTSD Checklist for *DSM-5* (PCL-5) with Life Events Checklist for *DSM-5* (LEC-5) and Criterion A

Version date: 11 April 2018

Reference: Weathers, F. W., Litz, B. T., Keane, T. M., Palmieri, P. A., Marx, B. P., & Schnurr, P. P. (2013). *The PTSD Checklist for DSM-5 (PCL-5) – LEC-5 and Extended Criterion A* [Measurement instrument]. Available from <https://www.ptsd.va.gov/>

URL: <https://www.ptsd.va.gov/professional/assessment/adult-sr/ptsd-checklist.asp>

Note: This is a fillable form. You may complete it electronically.

This page intentionally left blank

PCL-5 with LEC-5 and Criterion A

Part 1

Instructions: Listed below are a number of difficult or stressful things that sometimes happen to people. For each event check one or more of the boxes to the right to indicate that: (a) it happened to you personally; (b) you witnessed it happen to someone else; (c) you learned about it happening to a close family member or close friend; (d) you were exposed to it as part of your job (for example, paramedic, police, military, or other first responder); (e) you're not sure if it fits; or (f) it doesn't apply to you.

Be sure to consider your entire life (growing up as well as adulthood) as you go through the list of events.

Event	Happened to me	Witnessed it	Learned about it	Part of my job	Not sure	Doesn't apply
1. Natural disaster (for example, flood, hurricane, tornado, earthquake)						
2. Fire or explosion						
3. Transportation accident (for example, car accident, boat accident, train wreck, plane crash)						
4. Serious accident at work, home, or during recreational activity						
5. Exposure to toxic substance (for example, dangerous chemicals, radiation)						
6. Physical assault (for example, being attacked, hit, slapped, kicked, beaten up)						
7. Assault with a weapon (for example, being shot, stabbed, threatened with a knife, gun, bomb)						
8. Sexual assault (rape, attempted rape, made to perform any type of sexual act through force or threat of harm)						
9. Other unwanted or uncomfortable sexual experience						
10. Combat or exposure to a war-zone (in the military or as a civilian)						
11. Captivity (for example, being kidnapped, abducted, held hostage, prisoner of war)						
12. Life-threatening illness or injury						
13. Severe human suffering						
14. Sudden violent death (for example, homicide, suicide)						
15. Sudden accidental death						
16. Serious injury, harm, or death you caused to someone else						
17. Any other very stressful event or experience						

Part 2

A. If you checked anything for #17 in PART 1, briefly identify the event you were thinking of:

B. If you have experienced more than one of the events in PART 1, think about the event you consider the worst event, which for this questionnaire means the event that currently bothers you the most. If you have experienced only one of the events in PART 1, use that one as the worst event. Please answer the following questions about the worst event (check all options that apply):

Briefly describe the worst event (for example, what happened, who was involved, etc.).

How long ago did it happen? _____ (please estimate if you are not sure)

How did you experience it?

It happened to me directly

I witnessed it

I learned about it happening to a close family member or close friend

I was repeatedly exposed to details about it as part of my job (for example, paramedic, police, military, or other first responder)

Other, please describe _____

Was someone's life in danger?

Yes, my life

Yes, someone else's life

No

Was someone seriously injured or killed?

Yes, I was seriously injured

Yes, someone else was seriously injured or killed

No

Did it involve sexual violence? Yes No

If the event involved the death of a close family member or close friend, was it due to some kind of accident or violence, or was it due to natural causes?

Accident or violence

Natural causes

Not applicable (The event did not involve the death of a close family member or close friend)

How many times altogether have you experienced a similar event as stressful or nearly as stressful as the worst event?

Just once

More than once (please specify or estimate the total number of times you have had this experience _____)

Part 3

Below is a list of problems that people sometimes have in response to a very stressful experience. Keeping your worst event in mind, please read each problem carefully and then circle one of the numbers to the right to indicate how much you have been bothered by that problem in the past month.

In the past month, how much were you bothered by:	Not at all	A little bit	Moderately	Quite a bit	Extremely
1. Repeated, disturbing, and unwanted memories of the stressful experience?	0	1	2	3	4
2. Repeated, disturbing dreams of the stressful experience?	0	1	2	3	4
3. Suddenly feeling or acting as if the stressful experience were actually happening again (as if you were actually back there reliving it)?	0	1	2	3	4
4. Feeling very upset when something reminded you of the stressful experience?	0	1	2	3	4
5. Having strong physical reactions when something reminded you of the stressful experience (for example, heart pounding, trouble breathing, sweating)?	0	1	2	3	4
6. Avoiding memories, thoughts, or feelings related to the stressful experience?	0	1	2	3	4
7. Avoiding external reminders of the stressful experience (for example, people, places, conversations, activities, objects, or situations)?	0	1	2	3	4
8. Trouble remembering important parts of the stressful experience?	0	1	2	3	4
9. Having strong negative beliefs about yourself, other people, or the world (for example, having thoughts such as: I am bad, there is something seriously wrong with me, no one can be trusted, the world is completely dangerous)?	0	1	2	3	4
10. Blaming yourself or someone else for the stressful experience or what happened after it?	0	1	2	3	4
11. Having strong negative feelings such as fear, horror, anger, guilt, or shame?	0	1	2	3	4
12. Loss of interest in activities that you used to enjoy?	0	1	2	3	4
13. Feeling distant or cut off from other people?	0	1	2	3	4
14. Trouble experiencing positive feelings (for example, being unable to feel happiness or have loving feelings for people close to you)?	0	1	2	3	4
15. Irritable behavior, angry outbursts, or acting aggressively?	0	1	2	3	4
16. Taking too many risks or doing things that could cause you harm?	0	1	2	3	4
17. Being "superalert" or watchful or on guard?	0	1	2	3	4
18. Feeling jumpy or easily startled?	0	1	2	3	4
19. Having difficulty concentrating?	0	1	2	3	4
20. Trouble falling or staying asleep?	0	1	2	3	4

VA/DoD Clinical Practice Guidelines

Management of Posttraumatic Stress Disorder and Acute Stress Disorder



VA/DoD Evidence-Based Practice

Provider Summary

Version 4.0 | 2023



VA/DoD CLINICAL PRACTICE GUIDELINE FOR MANAGEMENT OF POSTTRAUMATIC STRESS DISORDER AND ACUTE STRESS DISORDER

Department of Veterans Affairs

Department of Defense

Provider Summary

QUALIFYING STATEMENTS

The Department of Veterans Affairs (VA) and the Department of Defense (DoD) guidelines are based on the best information available at the time of publication. The guidelines are designed to provide information and assist decision making. They are not intended to define a standard of care and should not be construed as one. Neither should they be interpreted as prescribing an exclusive course of management.

This clinical practice guideline (CPG) is based on a systematic review (SR) of both clinical and epidemiological evidence. Developed by a panel of multidisciplinary experts, it provides a clear explanation of the logical relationships between various care options and health outcomes while rating both the quality of the evidence and the strength of the recommendation.

Variations in practice will inevitably and appropriately occur when providers consider the needs of individual patients, available resources, and limitations unique to an institution or type of practice. Therefore, every health care professional using these guidelines is responsible for evaluating the appropriateness of applying them in the setting of any particular clinical situation with a patient-centered approach.

These guidelines are not intended to represent VA or DoD policies. Further, inclusion of recommendations for specific testing, therapeutic interventions, or both within these guidelines does not guarantee coverage of civilian sector care.

Version 4.0 - 2023

Table of Contents

Introduction	1
Highlights in this Guideline	2
Recommendations	3
Scope of the CPG	7
Guideline Development Team	7
Patient-centered Care	8
Shared Decision Making	8
Algorithm	9
Module A: Acute Stress Reaction/Disorder	10
Module B: Assessment and Diagnosis of Posttraumatic Stress Disorder	11
Module C: Management of Posttraumatic Stress Disorder	12
Pharmacotherapy Dosing Table	19
Methods	20
References	22

Introduction

The VA and DoD Evidence-Based Practice Work Group (EBPWG) was established and first chartered in 2004, with a mission to advise the VA/DoD Health Executive Committee “on the use of clinical and epidemiological evidence to improve the health of the population . . .” across the Veterans Health Administration (VHA) and Military Health System (MHS), by facilitating the development of CPGs for the VA and DoD populations.⁽¹⁾ Development and update of VA/DoD CPGs is funded by VA Evidence Based Practice, Office of Quality and Patient Safety. The system-wide goal of evidence-based CPGs is to improve patient health and wellbeing.

In 2017, VA and DoD published a CPG for Management of Posttraumatic Stress Disorder and Acute Stress Disorder (2017 VA/DoD PTSD CPG), which was based on evidence reviewed through March 2016. Since the release of that CPG, the evidence base on PTSD has expanded. Consequently, the EBPWG initiated the update of the 2017 VA/DoD PTSD CPG in 2022. This updated CPG’s use of Grading of Recommendations Assessment, Development, and Evaluation (GRADE) approach reflects a more rigorous application of the methodology than previous iterations.⁽²⁾ Therefore, the strength of some recommendations might have been modified because of the confidence in the quality of the supporting evidence (see Evidence Quality and Recommendation Strength in the full VA/DoD PTSD CPG).

This CPG provides an evidence-based framework for evaluating and managing care for individuals with posttraumatic stress disorder (PTSD) or acute stress disorder (ASD) toward improving clinical outcomes. Successful implementation of this CPG will:

- Assess the patient’s condition and collaborate with the patient, family, and caregivers to determine optimal management of patient care;
- Emphasize the use of patient-centered care and shared decision making;
- Minimize preventable complications and morbidity; and
- Optimize individual health outcomes and quality of life (QoL).

Highlights in this Guideline

The 2023 VA/DoD PTSD CPG reflects a more rigorous application of the GRADE methodology than the 2017 VA/DoD PTSD CPG. This approach has resulted in the exclusion or downgrading of studies included in previous versions of this CPG and has impacted the strength of some recommendations (e.g., *Strong for* downgraded to *Weak for*), despite a similar evidence base. For additional information on GRADE or CPG methodology, see Appendix A in the full CPG.

In the 2017 VA/DoD PTSD CPG, trauma-focused psychotherapies were evaluated as a class that included Cognitive Processing Therapy (CPT), Eye Movement Desensitization and Reprocessing (EMDR), and Prolonged Exposure (PE). In the 2023 CPG, the evidence on trauma-focused psychotherapies was reviewed for each treatment individually, rather than as a class, for comparison with medications. This review, combined with the more rigorous application of GRADE and accumulated new evidence, resulted in the downgrading of some specific treatments.

Significant changes to the strength of the recommendations include the following.

- Downgrading of some trauma-focused psychotherapies that previously received a *Strong for* recommendation (Brief Eclectic Psychotherapy and Narrative Exposure Therapy [NET] were downgraded to *Neither for nor against* [Recommendation 10], and Ehlers' Cognitive Therapy [CT] for PTSD and written narrative exposure, now called Written Exposure Therapy [WET], was downgraded to *Weak for* [Recommendation 9]).
- Downgrading of some non-trauma-focused psychotherapies that previously received a *Weak for* recommendation (Stress Inoculation Training [SIT] and Interpersonal Psychotherapy [IPT] were downgraded to *Neither for nor against* [Recommendation 10]).
- Downgrading of one medication that previously received a *Strong for* recommendation (Fluoxetine was downgraded to *Neither for nor against* [Recommendation 16]).
- Downgrading of some medications that previously received a *Weak for* recommendation (Nefazodone, phenelzine, and imipramine were downgraded to *Neither for nor against* [Recommendation 16]).
- Adding *Weak for* recommendations for prazosin for nightmares (Recommendation 32) and Mindfulness-Based Stress Reduction® (MBSR) for overall PTSD symptoms (Recommendation 26).

Other changes in the CPG include the following.

- Updated algorithm for screening and treatment of ASD and PTSD
- Greater attention to discussing the generalizability of evidence to subgroups based on gender identity, sexuality, race, ethnicity, age, and other patient characteristics

and clearer delineation of complementary, integrative, and alternative health treatment

- Updated recommendations on research needed to strengthen future guidelines

Recommendations

The evidence-based clinical practice recommendations listed (see [Table 1](#)) were made using a systematic approach considering four domains as per the GRADE approach (see Summary of Guideline Development Methodology in the full VA/DoD CPG). These domains include confidence in the quality of the evidence, balance of desirable and undesirable outcomes (i.e., benefits and harms), patient values and preferences, and other implications (e.g., resource use, equity, acceptability).

Table 1. Evidence-based Clinical Practice Recommendations with Strength and Category

Topic	Sub-topic	#	Recommendation	Strength ^a	Category ^b
Assessment and Diagnosis of PTSD		1.	When screening for PTSD, we suggest using the Primary Care PTSD Screen for DSM-5.	Weak for	Reviewed, New-replaced
		2.	For confirmation of the diagnosis of PTSD, we suggest using a validated structured clinician-administered interview, such as the Clinician-Administered PTSD Scale or PTSD Symptom Scale - Interview Version.	Weak for	Reviewed, New-replaced
		3.	To detect changes in PTSD symptom severity over time, we suggest the use of a validated instrument, such as the PTSD Checklist for DSM-5, or a structured clinician-administered interview, such as the Clinician-Administered PTSD Scale.	Weak for	Reviewed, New-replaced
Prevention of PTSD	<i>Selective Prevention of PTSD</i>	4.	For the prevention of PTSD among individuals who have been exposed to trauma, there is insufficient evidence to recommend for or against psychotherapy or pharmacotherapy in the immediate post-trauma period.	Neither for nor against	Not Reviewed, Amended
	<i>Indicated Prevention of PTSD</i>	5.	For the prevention of PTSD among patients diagnosed with acute stress disorder, we suggest trauma-focused cognitive behavioral psychotherapy.	Weak for	Reviewed, New-replaced
	<i>Indicated Prevention of PTSD</i>	6.	For the prevention of PTSD among patients diagnosed with acute stress reaction/acute stress disorder, there is insufficient evidence to recommend for or against any pharmacotherapy.	Neither for nor against	Reviewed, New-replaced

Topic	Sub-topic	#	Recommendation	Strength ^a	Category ^b
Treatment of PTSD	Treatment Selection	7.	We recommend individual psychotherapies, listed in Recommendation 8 , over pharmacologic interventions for the treatment of PTSD.	Strong for	Reviewed, New-replaced
	Psychotherapy	8.	We recommend the individual, manualized trauma-focused psychotherapies for the treatment of PTSD: Cognitive Processing Therapy, Eye Movement Desensitization and Reprocessing, or Prolonged Exposure.	Strong for	Reviewed, New-replaced
		9.	We suggest the following individual, manualized psychotherapies for the treatment of PTSD: Ehlers' Cognitive Therapy for PTSD, Present-Centered Therapy, or Written Exposure Therapy.	Weak for	Reviewed, New-replaced
		10.	There is insufficient evidence to recommend for or against the following individual psychotherapies for the treatment of PTSD: Accelerated Resolution Therapy, Adaptive Disclosure, Acceptance and Commitment Therapy, Brief Eclectic Psychotherapy, Dialectical Behavior Therapy, Emotional Freedom Techniques, Impact on Killing, Interpersonal Psychotherapy, Narrative Exposure Therapy, Prolonged Exposure in Primary Care, psychodynamic therapy, psychoeducation, Reconsolidation of Traumatic Memories, Seeking Safety, Stress Inoculation Training, Skills Training in Affective and Interpersonal Regulation, Skills Training in Affective and Interpersonal Regulation in Primary Care, supportive counseling, Thought Field Therapy, Trauma-Informed Guilt Reduction, or Trauma Management Therapy.	Neither for nor against	Reviewed, New-replaced
		11.	There is insufficient evidence to recommend using individual components of manualized psychotherapy protocols over, or in addition to, the full therapy protocol for the treatment of PTSD.	Neither for nor against	Reviewed, Not Changed
		12.	There is insufficient evidence to recommend for or against any specific manualized group therapy for the treatment of PTSD.	Neither for nor against	Reviewed, New-replaced
		13.	There is insufficient evidence to recommend using group therapy as an adjunct for the primary treatment of PTSD.	Neither for nor against	Reviewed, New-replaced
		14.	There is insufficient evidence to recommend for or against the following couples therapies for the treatment of PTSD: Behavioral Family Therapy, Structured Approach Therapy, or Cognitive Behavioral Conjoint Therapy.	Neither for nor against	Reviewed, Not Changed

Topic	Sub-topic	#	Recommendation	Strength ^a	Category ^b
Treatment of PTSD (cont.)	Pharmacotherapy	15.	We recommend paroxetine, sertraline, or venlafaxine for the treatment of PTSD.	Strong for	Reviewed, New-replaced
		16.	There is insufficient evidence to recommend for or against amitriptyline, bupropion, buspirone, citalopram, desvenlafaxine, duloxetine, escitalopram, eszopiclone, fluoxetine, imipramine, mirtazapine, lamotrigine, nefazodone, olanzapine, phenelzine, pregabalin, rivastigmine, topiramate, or quetiapine for the treatment of PTSD.	Neither for nor against	Reviewed, New-replaced
	Pharmacotherapy (cont.)	17.	There is insufficient evidence to recommend for or against psilocybin, ayahuasca, dimethyltryptamine, ibogaine, or lysergic acid diethylamide for the treatment of PTSD.	Neither for nor against	Reviewed, New-added
		18.	We suggest against divalproex, guanfacine, ketamine, prazosin, risperidone, tiagabine, or vortioxetine for the treatment of PTSD.	Weak against	Reviewed, New-replaced
		19.	We recommend against benzodiazepines for the treatment of PTSD.	Strong against	Reviewed, New-replaced
		20.	We recommend against cannabis or cannabis derivatives for the treatment of PTSD.	Strong against	Reviewed, Amended
	Augmentation Therapy	21.	There is insufficient evidence to recommend for or against the combination or augmentation of psychotherapy (see Recommendation 8 and Recommendation 9) or medications (see Recommendation 15) with any psychotherapy or medication for the treatment of PTSD (see Recommendation 22 for antipsychotic medications and Recommendation 23 for 3,4-methylenedioxymethamphetamine).	Neither for nor against	Reviewed, New-replaced
		22.	We suggest against aripiprazole, asenapine, brexpiprazole, cariprazine, iloperidone, lumateperone, lurasidone, olanzapine, paliperidone, quetiapine, risperidone, or ziprasidone for augmentation of medications for the treatment of PTSD.	Weak against	Reviewed, New-replaced
		23.	There is insufficient evidence to recommend for or against 3,4-methylenedioxymethamphetamine assisted psychotherapy for the treatment of PTSD.	Neither for nor against	Reviewed, New-added
	Non-pharmacologic Biological Treatments	24.	There is insufficient evidence to recommend for or against the following somatic therapies for the treatment of PTSD: capnometry-assisted respiratory therapy, hyperbaric oxygen therapy, neurofeedback, NightWare®, repetitive transcranial magnetic stimulation, stellate ganglion block, or transcranial direct current stimulation.	Neither for nor against	Reviewed, New-replaced
		25.	We suggest against electroconvulsive therapy or vagus nerve stimulation for treatment of PTSD.	Weak against	Reviewed, New-replaced
		26.	We suggest Mindfulness-Based Stress Reduction® for the treatment of PTSD.	Weak for	Reviewed, New-replaced

Topic	Sub-topic	#	Recommendation	Strength ^a	Category ^b
Treatment of PTSD (cont.)	Complementary, Integrative, and Alternative Approaches	27.	There is insufficient evidence to recommend for or against the following mind-body interventions for the treatment of PTSD: acupuncture, Cognitively Based Compassion Training Veteran version, creative arts therapies (e.g., music, art, dance), guided imagery, hypnosis or self-hypnosis, Loving Kindness Meditation, Mantram Repetition Program, Mindfulness-Based Cognitive Therapy, other mindfulness trainings (e.g., integrative exercise, Mindfulness-Based Exposure Therapy, brief mindfulness training), relaxation training, somatic experiencing, tai chi or qigong, Transcendental Meditation®, and yoga.	Neither for nor against	Reviewed, New-replaced
		28.	There is insufficient evidence to recommend for or against the following interventions for the treatment of PTSD: recreational therapy, aerobic or non-aerobic exercise, animal-assisted therapy (e.g., canine, equine), and nature experiences (e.g., fishing, sailing).	Neither for nor against	Reviewed, New-replaced
	Technology-based Treatment	29.	We recommend secure video conferencing to deliver treatments in Recommendation 8 and Recommendation 9 when that therapy has been validated for use with video conferencing or when other options are unavailable.	Strong for	Reviewed, New-replaced
		30.	There is insufficient evidence to recommend for or against mobile apps or other self-help-based interventions for the treatment of PTSD.	Neither for nor against	Reviewed, New-added
		31.	There is insufficient evidence to recommend for or against facilitated internet-based cognitive behavioral therapy for the treatment of PTSD.	Neither for nor against	Reviewed, New-replaced
Treatment of Nightmares		32.	We suggest prazosin for the treatment of nightmares associated with PTSD.	Weak for	Reviewed, Amended
		33.	There is insufficient evidence to recommend for or against the following treatments for nightmares associated with PTSD: Imagery Rehearsal Therapy, Exposure Relaxation and Rescripting Therapy, Imaging Rescripting and Reprocessing Therapy, or NightWare.	Neither for nor against	Reviewed, New-added
Treatment of PTSD with Co-Occurring Conditions		34.	We suggest that the presence of co-occurring substance use disorder and/or other disorder(s) not preclude treatments in Recommendation 8 and Recommendation 9 for PTSD.	Weak for	Reviewed, New-replaced

^a For additional information, see Grading Recommendations in the full VA/DoD PTSD CPG.

^b For additional information, see Recommendation Categorization and Appendix E in the full VA/DoD PTSD CPG.

Scope of the CPG

This CPG is based on published clinical evidence and related information available through May 4, 2022. It is intended to provide general guidance on best evidence-based practices (see Appendix A in the full VA/DoD PTSD CPG for additional information on the evidence review methodology). Although the CPG is intended to improve the quality of care and clinical outcomes (see Introduction in the full VA/DoD PTSD CPG), it is not intended to define a standard of care (i.e., mandated or strictly required care).

This CPG is intended for use by providers and others involved in the care of active duty Service members and Veterans with PTSD.

The patient population of interest for this CPG is adults with PTSD or ASD caused by any type of trauma who are eligible for care in VA or DoD health care delivery systems.

Guideline Development Team

Table 2. Guideline Work Group and Guideline Development Team

Organization	Name*
Department of Veterans Affairs	Paula P. Schnurr, PhD
	Jessica L. Hamblen, PhD
	Claire Collie, PhD
	Matthew A. Fuller, PharmD, FASHP, BCPP
	Paul E. Holtzheimer, MD, MS
	Ursula Kelly, PhD, APRN, ANP-BC, PMHNP-BC, FAANP, FAAN
	Ariel Lang, PhD, MPH
	Sonya Norman, PhD
	Katie Papke, LMSW, CAADC, CCTP, CHTVSP
	Ismene Petrakis, MD
	Brian Shiner, MD, MPH
	Ilse Wiechers, MD, MPP, MHS
Department of Defense	Marija Kelber, PhD
	Jonathan Wolf, MD
	Rachael Coller, PharmD
	Kate McGraw, PhD
	CAPT Joshua Morganstein, MD
David Riggs, PhD	
VA Evidence Based Practice, Office of Quality and Patient Safety Veterans Health Administration	James Sall, PhD, FNP-BC
	René Sutton, BS, HCA
	Jennifer Ballard-Hernandez, DNP, RN, FNP-BC

Organization	Name*
Clinical Quality Improvement Program Defense Health Agency	Elaine P. Stuffel, MHA, BSN, RN
	Cynthia F. Villareal, BSN, RN
	Isabella Alvarez, MA, BSN, RN
The Lewin Group	Clifford Goodman, PhD
	Jennifer Weil, PhD
	Peter Baroff, BA
	Charlie Zachariades, MSc
	Annie Zhang, BA
ECRI	Jim Reston, PhD, MPH
	Amy Tsou, MD, MSc
	Rebecca Rishar, BA, MLIS
Sigma Health Consulting	Frances M. Murphy, MD, MPH
	James G. Smirniotopoulos, MD
Duty First Consulting	Kate Johnson, BS
	Rachel Piccolino, BA

*Additional contributor contact information is available in Appendix F in the full VA/DoD PTSD CPG.

Patient-centered Care

Intended to consider patient needs and preferences, guideline recommendations represent a whole/holistic health approach to care that is patient centered, culturally appropriate, and available to people with limited literacy skills and physical, sensory, or learning disabilities. VA/DoD CPGs encourage providers to use a patient-centered, whole/holistic health approach (i.e., individualized treatment based on patient needs, characteristics, and preferences). This approach aims to treat the particular condition while also optimizing the individual’s overall health and wellbeing.

Regardless of the care setting, all patients should have access to individualized evidence-based care. Patient-centered care can decrease patient anxiety, increase trust in providers, and improve treatment adherence.(3, 4) A whole/holistic health approach (<https://www.va.gov/wholehealth/>) empowers and equips individuals to meet their personal health and wellbeing goals. Good communication is essential and should be supported by evidence-based information tailored to each patient’s needs. An empathetic and non-judgmental approach facilitates discussions sensitive to gender, culture, ethnicity, and other differences.

Shared Decision Making

This CPG encourages providers to practice shared decision making, a process in which providers, patients, and patient care partners (e.g., family, friends, caregivers) consider clinical evidence of benefits and risks as well as patient values and preferences to make





decisions regarding the patient’s treatment.⁽⁵⁾ Shared decision making is emphasized in *Crossing the Quality Chasm*, an Institute of Medicine, now NAM, report in 2001 ⁽⁶⁾ and is inherent within the whole/holistic health approach. Providers must be adept at presenting information to their patients regarding individual treatments, expected risks, expected outcomes, and levels or settings of care or both, especially where patient heterogeneity in weighing risks and benefits might exist. VHA and MHS have embraced shared decision making. Providers are encouraged to use shared decision making to individualize treatment goals and plans based on patient capabilities, needs, and preferences.

Algorithm

This CPG’s algorithm is designed to facilitate understanding of the clinical pathway and decision-making process used in managing patients with PTSD. This algorithm format represents a simplified flow of the management of patients with PTSD and helps foster efficient decision making by providers. It includes


- Steps of care in an ordered sequence,
- Decisions to be considered,
- Decision criteria recommended, and
- Actions to be taken.

The algorithm is a step-by-step decision tree. Standardized symbols display each step, and arrows connect the numbered boxes indicating the order in which the steps should be followed.⁽⁷⁾ Sidebars 1–11 provide more detailed information to assist in defining and interpreting elements in the boxes.

Shape	Description
	Rounded rectangles represent a clinical state or condition.
	Hexagons represent a decision point in the process of care, formulated as a question that can be answered “Yes” or “No.”
	Rectangles represent an action in the process of care.
	Ovals represent a link to another section within the algorithm.


For alternative text descriptions of the algorithm, please refer to Appendix H in the full VA/DoD CPG PTSD.

Module A: Acute Stress Reaction/Disorder




Abbreviations: ASR: Acute Stress Reaction; COSR: Combat and Operational Stress Reaction; CPG: clinical practice guideline; DSM: Diagnostic and Statistical Manual of Mental Disorders; PTSD: Posttraumatic stress disorder; VA: Veteran Affairs; DoD: Department of Defense.

Module B: Assessment and Diagnosis of Posttraumatic Stress Disorder



Abbreviations: CPG: clinical practice guideline; PTSD: Posttraumatic stress disorder; VA: Veteran Affairs; DoD: Department of Defense.

Module C: Management of Posttraumatic Stress Disorder



Abbreviations: CPG: clinical practice guideline; PTSD: Posttraumatic stress disorder; VA: Veteran Affairs; DoD: Department of Defense.

Sidebar 1: Immediate Needs

- Survival (including first aid and stabilizing physical condition), safety, and security
- Food, hydration, shelter, and clothing
- Sleep
- Orientation
- Communication with unit, family, friends, and community
- Education and normalization of reactions to trauma

Sidebar 2: Assessment

- History of trauma and mental health concerns
- Symptoms
- Consider screening for PTSD symptoms using the PC-PTSD-5 (Recommendation 1)
- Medical status
- Mental status, including suicidality (consult VA/DoD CPG for Assessment and Management of Patients at Risk for Suicide,^a as needed)
- Functional status
- Psychosocial status, including intimate and family relationships; financial problems; legal issues
- Occupational performance
- Substance use
- Strengths, coping skills, and protective factors

^a See the VA/DoD CPG for the Assessment and Management of Patients at Risk for Suicide, available at <https://www.healthquality.va.gov/>.

Abbreviations: PC-PTSD-5: Primary Care PTSD Screen for Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition; PTSD: posttraumatic stress disorder; VA: Veteran Affairs; DoD: Department of Defense.

Sidebar 3: DSM-5-TR Diagnostic Criteria for Acute Stress Disorder(8)

Criterion A. Exposure to actual or threatened death, serious injury, or sexual violence in one (or more) of the following ways:

1. Directly experiencing the traumatic event(s).
2. Witnessing, in person, the event(s) as it occurred to others.
3. Learning that the event(s) occurred to a close family member or close friend. **Note:** In cases of actual or threatened death of a family member or friend, the event(s) must have been violent or accidental.
4. Experiencing repeated or extreme exposure to aversive details of the traumatic event(s) (e.g., first responders collecting human remains, police officers repeatedly exposed to details of child abuse).

Note: This does not apply to exposure through electronic media, television, movies, or pictures unless this exposure is work-related.

Sidebar 3: DSM-5-TR Diagnostic Criteria for Acute Stress Disorder(8)

Criterion B. Presence of nine (or more) of the following symptoms from any of the five categories of intrusion, negative mood, dissociation, avoidance, and arousal, beginning or worsening after the traumatic event(s) occurred:

Intrusion Symptoms

1. Recurrent, involuntary, and intrusive distressing memories of the traumatic event(s).
2. Recurrent distressing dreams in which the content and/or affect of the dream are related to the traumatic event(s).
3. Dissociative reactions (e.g., flashbacks) in which the individual feels or acts as if the traumatic event(s) were recurring. (Such reactions may occur on a continuum, with the most extreme expression being a complete loss of awareness of present surroundings.)
4. Intense or prolonged psychological distress or marked physiological reactions in response to internal or external cues that symbolize or resemble an aspect of the traumatic event(s).

Negative Mood

5. Persistent inability to experience positive emotions (e.g., inability to experience happiness, satisfaction, or loving feelings).

Dissociative Symptoms

6. An altered sense of the reality of one's surroundings or oneself (e.g., seeing oneself from another's perspective, being in a daze, time slowing).
7. Inability to remember an important aspect of the event(s) (typically due to dissociative amnesia and not to other factors such as head injury, alcohol, or drugs).

Avoidance Symptoms

8. Efforts to avoid distressing memories, thoughts, or feelings about or closely associated with the traumatic event(s).
9. Efforts to avoid external reminders (people, places, conversations, activities, objects, situations) that arouse distressing memories, thoughts, or feelings about or closely associated with the traumatic event(s).

Arousal Symptoms

10. Sleep disturbance (e.g., difficulty falling or staying asleep, restless sleep).
11. Irritable behavior and angry outbursts (with little or no provocation), typically expressed as verbal or physical aggression toward people or objects.
12. Hypervigilance.
13. Problems with concentration.
14. Exaggerated startle response.

Criterion C. Duration of the disturbance (symptoms in Criterion B) is 3 days to 1 month after trauma exposure.

Note: Symptoms typically begin immediately after the trauma, but persistence for at least 3 days and up to 1 month is needed to meet disorder criteria.

Criterion D. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.

Criterion E. The disturbance is not attributable to the physiological effects of a substance (e.g., medication or alcohol) or another medical condition (e.g., mild traumatic brain injury) and is not better explained by a brief psychotic disorder.

Sidebar 4: Acute Interventions for Acute Stress Disorder

- Individual, manualized trauma-focused cognitive behavioral psychotherapy
- Consider: Collaborative care or wellness-oriented activities

Sidebar 5: Acute Interventions for Acute Stress Response/Combat and Operational Stress Reaction

- Education and normalization, acute symptom management, social support
- Suggest: Brief cognitive behavioral psychotherapy

Sidebar 6: General Assessment

- Complete comprehensive clinical assessment of presenting complaints and comorbid conditions
- Perform safety, lethal means, and environmental assessment
- Consider history and presenting complaints: mental health, medical, military, marital, family, substance use, social and spiritual life, functional status
- Identify lifetime trauma history and duration of exposure
- Record current and past medications (including over-the-counter drugs and herbals) and psychosocial treatment
- Consider, with patient consent, obtaining an additional history from family, significant other, or both
- Perform mental status exam
- Consider, in cases of diagnostic uncertainty, use of validated structured clinical interviews for PTSD (i.e., CAPS-5, PSSI) (see Recommendation 2)

Abbreviations: CAPS-5: Clinician-Administered Posttraumatic Stress Disorder Scale for Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, Text Revision; PSSI: PTSD Symptom Scale - Interview Version

Sidebar 7: DSM-5-TR Diagnostic Criteria for PTSD(8)

Criterion A. Exposure to actual or threatened death, serious injury, or sexual violence in one (or more) of the following ways:

1. Directly experiencing the traumatic event(s).
2. Witnessing, in person, the event(s) as it occurred to others.
3. Learning that the traumatic event(s) occurred to a close family member or close friend. In cases of actual or threatened death of a family member or friend, the event(s) must have been violent or accidental.
4. Experiencing repeated or extreme exposure to aversive details of the traumatic event(s) (e.g., first responders collecting human remains, police officers repeatedly exposed to details of child abuse).

Note: Criterion A4 does not apply to exposure through electronic media, television, movies, or pictures unless this exposure is work related.

Sidebar 7: DSM-5-TR Diagnostic Criteria for PTSD(8)

Criterion B. Presence of one (or more) of the following intrusion symptoms associated with the traumatic event(s), beginning after the traumatic event(s) occurred:

1. Recurrent, involuntary, and intrusive distressing memories of the traumatic event(s).
2. Recurrent distressing dreams in which the content and/or affect of the dream are related to the traumatic event(s).
3. Dissociative reactions (e.g., flashbacks) in which the individual feels or acts as if the traumatic event(s) were recurring. (Such reactions may occur on a continuum, with the most extreme expression being a complete loss of awareness of present surroundings)
4. Intense or prolonged psychological distress at exposure to internal or external cues that symbolize or resemble an aspect of the traumatic event(s).
5. Marked physiological reactions to internal or external cues that symbolize or resemble an aspect of the traumatic event(s).

Criterion C. Persistent avoidance of stimuli associated with the traumatic event(s), beginning after the traumatic event(s) occurred, as evidenced by one or both of the following:

1. Avoidance of or efforts to avoid distressing memories, thoughts, or feelings about or closely associated with the traumatic event(s).
2. Avoidance or efforts to avoid external reminders (people, places, conversations, activities, objects, situations) that arouse distressing memories, thoughts, or feelings about or closely associated with the traumatic event(s).

Criterion D. Negative alterations in cognitions and mood associated with the traumatic event(s), beginning or worsening after the traumatic event(s) occurred, as evidenced by two (or more) of the following:

1. Inability to recall an important aspect of the traumatic event(s) (typically due to dissociative amnesia and not to other factors such as head injury, alcohol, or drugs).
2. Persistent and exaggerated negative beliefs or expectations about oneself, others, or the world (e.g., “I am bad,” “No one can be trusted,” “The world is completely dangerous,” “My whole nervous system is permanently ruined”).
3. Persistent, distorted cognitions about the cause or consequences of the traumatic event(s) that lead the individual to blame themselves or others.
4. Persistent negative emotional state (e.g., fear, horror, anger, guilt, shame).
5. Markedly diminished interest or participation in significant activities.
6. Feeling of detachment or estrangement from others.
7. Persistent inability to experience positive emotions (e.g., inability to experience happiness, satisfaction, or loving feelings).

Criterion E. Marked alterations in arousal and reactivity associated with the traumatic event(s), beginning or worsening after the traumatic event(s) occurred, as evidenced by two (or more) of the following:

1. Irritable behavior and angry outbursts (with little or no provocation) typically expressed as verbal or physical aggression toward people or objects.
2. Reckless or self-destructive behavior.
3. Hypervigilance.
4. Exaggerated startle response.
5. Problems with concentration.
6. Sleep disturbance (e.g., difficulty falling or staying asleep or restless sleep).

Criterion F. Duration of the disturbance (Criteria B, C, D, and E) is more than 1 month.

Sidebar 7: DSM-5-TR Diagnostic Criteria for PTSD(8)

Criterion G. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.

Criterion H. The disturbance is not attributable to the physiological effects of a substance (e.g., medication, alcohol) or another medical condition.

Specify whether:

With dissociative symptoms: The individual's symptoms meet the criteria for posttraumatic stress disorder, and in addition, in response to the stressor, the individual experiences persistent or recurrent symptoms of either of the following:

1. **Depersonalization:** Persistent or recurrent experiences of feeling detached from, and as if one were an outside observer of, one's mental processes or body (e.g., feeling as though one were in a dream; feeling a sense of unreality of self or body or of time moving slowly).
2. **Derealization:** Persistent or recurrent experiences of unreality of surroundings (e.g., the world around the individual is experienced as unreal, dreamlike, distant, or distorted).

Note: To use this subtype, the dissociative symptoms must not be attributable to the physiological effects of a substance (e.g., blackouts, behavior during alcohol intoxication) or another medical condition (e.g., complex partial seizures).

Specify if:

With delayed expression: If the full diagnostic criteria are not met until at least 6 months after the event (although the onset and expression of some symptoms may be immediate).

Sidebar 8: Assessment of PTSD Symptoms

- Assess PTSD symptoms using validated instruments, such as the PTSD Checklist for DSM-5 (PCL-5), or a structured clinician-administered interview (e.g., CAPS-5) (see Recommendation 3).

Abbreviations: CAPS-5: Clinician-Administered PTSD Scale for DSM-5; DSM-5: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition; PCL-5: PTSD Checklist for DSM-5; PTSD: posttraumatic stress disorder

Sidebar 9: Treatment Selection

1. Initiate recommended individual, manualized psychotherapy (see Recommendation 8) according to patient preference.
2. If individual psychotherapy is unavailable or not preferred, initiate recommended pharmacotherapy (see Recommendation 15).
3. If options 1 and 2 are infeasible or are not preferred, offer suggested psychotherapy (see Recommendation 9) or suggested CIH (see Recommendation 26).
4. If options 1, 2, and 3 are infeasible or are not preferred, consider other psychotherapies (see Recommendation 10, Recommendation 12, and Recommendation 14), other pharmacotherapy (see Recommendation 16), complementary, integrative, or alternative approaches (see Recommendation 27 and Recommendation 28) based on availability, patient preference, and review of current evidence.
5. If none of the options above are acceptable to the patient, consider treating other disorders, issues, or both and reevaluating for PTSD treatment later.

Abbreviations: CIH: complementary and integrative health; PTSD: posttraumatic stress disorder

Sidebar 10: Additional Treatment and Support Needs

- Consider treatment for comorbidities and other identified problems (see Recommendation 34 as well as other relevant VA/DoD CPGs*).
- Consider symptom-specific management (e.g., sleep, pain).
- Facilitate social support.
- Address Whole Health by offering CIH, alternative approaches, health and wellbeing coaching, recreation therapy, etc.

*VA/DoD CPGs can be found at the following link: <https://www.healthquality.va.gov/index.asp>. Relevant VA/DoD CPGs to consult might include CPGs for the Management of Major Depressive Disorder, Substance Use Disorder, Bipolar Disorder, Suicide, Chronic Multisymptomatic Illness, Concussion-Mild Traumatic Brain Injury, and others.

Abbreviations: CIH: complementary and integrative health

Sidebar 11: Clinically Meaningful Response Time

- Psychotherapies require an adequate dosage to be fully effective in reducing PTSD symptoms; some effects might also not become apparent until some period has elapsed after treatment is initiated. For the indicated psychotherapies for PTSD (see Recommendation 8), it is generally accepted that initial treatment effects will be noticeable after 4–8 sessions typically delivered over 8–12 weeks. Psychotherapies might have an attenuated effect if delivered less than weekly.
- The pharmacological management of PTSD requires the SSRI (e.g., sertraline) or SNRI (e.g., venlafaxine) be given at an appropriate dosage for an adequate time to allow for the full therapeutic effects before moving to alternative or augmentative treatment options. These medications should be initiated at the recommended starting dose and titrated based on clinical response and tolerability (see Appendix B in the full VA/DoD PTSD CPG). The duration of the trial should be 8–12 weeks.

Abbreviations: PTSD: posttraumatic stress disorder; SNRI: serotonin-norepinephrine reuptake inhibitor; SSRI: selective serotonin reuptake inhibitor

Sidebar 12: Maintenance Plan

- Terminate PTSD treatment or taper based on clinician judgment and patient preference, normalize fluctuations in symptoms, discuss self-monitoring for symptoms that warrant future attention, and provide resources for seeking care in the future.
- Before termination of psychopharmacology, discuss the risks and benefits of discontinuing medication, including possible side effects and return of symptoms. Make a schedule to taper based on patient preference with a discussion of the length of time required and consideration of anticipated life events and stressors. Discuss the plan for monitoring during and post taper, including steps needed to reinstate pharmacology.
- Should the patient wish to continue pharmacotherapy, investigate, and discuss continuing medications with behavioral health or primary care.
- Refer the patient for treatment of other disorders or functional issues (e.g., relationship distress).
- If desired, facilitate referral to health and wellbeing programs as a part of a Whole Health approach to care.

Abbreviations: PTSD: posttraumatic stress disorder

Pharmacotherapy Dosing Table

Table 3: Pharmacotherapy Dosing Table

Therapeutic Category	Initial Dose	Dose Range	Clinical Considerations: Comorbidities and Safety
Antidepressants Monotherapy <ul style="list-style-type: none"> • Paroxetine • Sertraline • Venlafaxine 	IR: 10–20 mg daily CR: 12.5 mg daily 25–50 mg daily IR: 25 mg 2 or 3 times a day XR: 37.5 mg daily	20–50 mg daily 12.5–50 mg 50–200 mg daily 75–375 mg in 2–3 divided doses 75–300 mg once daily	<ul style="list-style-type: none"> • All antidepressants have a Black Box warning for increased risk of suicidality in children and young adults (≤24 years) in short-term studies of MDD and other psychiatric disorders. • Avoid abrupt discontinuation; withdrawal symptoms can occur with sudden discontinuation of SSRIs and SNRIs, paroxetine and venlafaxine in particular. • Paroxetine and sertraline have FDA indications for treating PTSD. • Common adverse effects of the SSRIs and SNRIs include nausea, headache, diarrhea, nervousness, sexual dysfunction, dizziness. • Hyponatremia or SIADH can occur; risk is elevated in patients >65 years. • Venlafaxine can elevate blood pressure; caution is advised with patients with hypertension. • Serotonin syndrome can occur, especially with concomitant medications that affect serotonin.
Other Agents <ul style="list-style-type: none"> • Prazosin# 	1 mg at bedtime; titrate to clinical response.	3–20 mg at bedtime	<ul style="list-style-type: none"> • Prazosin might cause syncope with a sudden loss of consciousness. Syncopal episodes usually occur within 30 to 90 minutes of the initial dose. • Addition of a diuretic, other antihypertensive agents, or a PDE5 inhibitor might cause an additive hypotensive effect.

Abbreviations: CR: controlled release; FDA: Food and Drug Administration; IR: immediate release; MDD: major depressive disorder; mg: milligram; PDE5: phosphodiesterase-5; PTSD: posttraumatic stress disorder; SIADH: syndrome of inappropriate antidiuretic hormone; SNRI: serotonin/norepinephrine reuptake inhibitor; SSRI: selective serotonin reuptake inhibitor; XR: extended release

for the treatment of nightmares associated with PTSD

Methods

The methodology used in developing this CPG follows the *Guideline for Guidelines*, an internal document of the VA/DoD EBPWG updated in January 2019 that outlines procedures for developing and submitting VA/DoD CPGs.⁽¹⁾ The *Guideline for Guidelines* is available at <http://www.healthquality.va.gov/policy/index.asp>. This CPG also aligns with the National Academy of Medicine’s (NAM) principles of trustworthy CPGs (e.g., explanation of evidence quality and strength, management of potential conflicts of interest [COI], interdisciplinary stakeholder involvement, use of SR and external review).⁽⁹⁾ Appendix A in the full VA/DoD PTSD CPG provides a detailed description of the CPG development methodology.

The Work Group used the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach to craft each recommendation and determine its strength. Per GRADE approach, recommendations must be evidence-based and cannot be made based on expert opinion alone. The GRADE approach uses the following four domains to inform the strength of each recommendation: confidence in the quality of the evidence, balance of desirable and undesirable outcomes, patient values and preferences, other considerations as appropriate (e.g., resource use, equity) (see Grading Recommendations in the full VA/DoD PTSD CPG).⁽¹⁰⁾

Using these four domains, the Work Group determined the relative strength of each recommendation (*Strong* or *Weak*). The strength of a recommendation is defined as the extent to which one can be confident that the desirable effects of an intervention outweigh its undesirable effects and is based on the framework above, which incorporates the four domains.⁽¹¹⁾ A *Strong* recommendation generally indicates *High* or *Moderate* confidence in the quality of the available evidence, a clear difference in magnitude between the benefits and harms of an intervention, similar patient values and preferences, and understood influence of other implications (e.g., resource use, feasibility).

In some instances, insufficient evidence exists on which to base a recommendation for or against a particular therapy, preventive measure, or other intervention. For example, the systematic evidence review might have found little or no relevant evidence, inconclusive evidence, or conflicting evidence for the intervention. The manner in which this finding is expressed in the CPG might vary. In such instances, the Work Group might include among its set of recommendations a statement of insufficient evidence for an intervention that might be in common practice although it is unsupported by clinical evidence and particularly if other risks of continuing its use might exist (e.g., high opportunity cost, misallocation of resources). In other cases, the Work Group might decide to exclude this type of statement about an intervention. For example, the Work Group might remain silent where an absence of evidence occurs for a rarely used intervention. In other cases, an intervention might have a favorable balance of benefits and harms but might be a standard of care for which no recent evidence has been generated.

Using these elements, the Work Group determines the strength and direction of each recommendation and formulates the recommendation with the general corresponding text as shown in Table 4.

Table 4. Strength and Direction of Recommendations and General Corresponding Text


Recommendation Strength and Direction	General Corresponding Text
Strong for	We recommend . . .
Weak for	We suggest . . .
Neither for nor against	There is insufficient evidence to recommend for or against . . .
Weak against	We suggest against . . .
Strong against	We recommend against . . .

The GRADE of each recommendation made in the 2023 CPG can be found in the section on [Recommendations](#). Additional information regarding the use of the GRADE system can be found in Appendix A in the full VA/DoD PTSD CPG.

References

1. U.S. Department of Veterans Affairs/Department of Defense Health Executive Committee (HEC). Evidence Based Practice Work Group Charter [updated January 9, 2017]. Available from: www.healthquality.va.gov/documents/EvidenceBasedPracticeWGCharter123020161.pdf.
2. Guyatt GH, Oxman AD, Kunz R, Atkins D, Brozek J, Vist G, et al. GRADE guidelines: 2. Framing the question and deciding on important outcomes. *Journal of clinical epidemiology*. 2011;64(4):395-400. Epub 2011/01/05. doi: 10.1016/j.jclinepi.2010.09.012. PubMed PMID: 21194891.
3. Robinson JH, Callister LC, Berry JA, Dearing KA. Patient-centered care and adherence: Definitions and applications to improve outcomes. *Journal of the American Academy of Nurse Practitioners*. 2008;20(12):600-7. Epub 2009/01/06. doi: 10.1111/j.1745-7599.2008.00360.x. PubMed PMID: 19120591.
4. Stewart M, Brown JB, Donner A, McWhinney IR, Oates J, Weston WW, et al. The impact of patient-centered care on outcomes. *J Fam Pract*. 2000;49(9):796-804. Epub 2000/10/14. PubMed PMID: 11032203.
5. National Learning Consortium. Shared Decision Making 2013. Available from: https://www.healthit.gov/sites/default/files/nlc_shared_decision_making_fact_sheet.pdf.
6. Institute of Medicine. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington DC: National Academies Press, 2001.
7. Society for Medical Decision Making Committee on Standardization of Clinical Algorithms. Proposal for clinical algorithm standards. *Med Decis Making*. 1992;12(2):149-54. Epub 1992/04/01. PubMed PMID: 1573982.
8. Association AP. *Diagnostic and statistical manual of mental disorders, Text Revision (DSM-5-TR®)*. Fifth, Text Revision ed. Arlington, VA: American Psychiatric Association; 2022.
9. Ransohoff DF, Pignone M, Sox HC. How to decide whether a clinical practice guideline is trustworthy. *Jama*. 2013;309(2):139-40. Epub 2013/01/10. doi: 10.1001/jama.2012.156703. PubMed PMID: 23299601.
10. Andrews JC, Schunemann HJ, Oxman AD, Pottie K, Meerpohl JJ, Coello PA, et al. GRADE guidelines: 15. Going from evidence to recommendation-determinants of a recommendation's direction and strength. *Journal of clinical epidemiology*. 2013;66(7):726-35. Epub 2013/04/11. doi: 10.1016/j.jclinepi.2013.02.003. PubMed PMID: 23570745.
11. Andrews J, Guyatt G, Oxman AD, Alderson P, Dahm P, Falck-Ytter Y, et al. GRADE guidelines: 14. Going from evidence to recommendations: the significance and presentation of recommendations. *Journal of clinical epidemiology*. 2013;66(7):719-25. Epub 2013/01/15. doi: 10.1016/j.jclinepi.2012.03.013. PubMed PMID: 23312392.

Access to the full guideline and additional resources is available at:
<https://www.healthquality.va.gov/>.



Helpful Prompts for evaluating PTSD Symptoms

(from the Clinician Administered PTSD Scale, Weathers, et al., 2013)

Criterion A: The index trauma

Definition	Clarifying Prompts
<p>The experience of trauma involves exposure to <u>actual</u> or <u>threatened death</u>, <u>serious injury</u>, or <u>sexual violence</u> that is:</p> <ul style="list-style-type: none"> • <u>Directly</u> experienced • <u>Witnessed</u> as it occurred to others • <u>Learned about</u> having occurred to a close family member or friend (must be violent or accidental) • Experienced as repeated or extreme exposure to aversive details of the event. • Not movies, television, pictures or other media unless <u>work related</u>. 	<p><i>What happened?</i> <i>How old were you?</i> <i>How were you involved?</i> <i>Who else was involved?</i> <i>Was anyone seriously injured or killed?</i> <i>Was anyone's life in danger?</i> <i>How many times did this happen?</i> <i>Did it involve sexual violence?</i></p> <p>If unclear, ask directly: <i>Is this something you experienced, [witnessed or learned about]?</i></p> <p>If it involved the death of a close friend or family member: <i>Was it due to some kind of accident or violence, or was it due to natural causes?</i></p> <p>With regard to exposure to repeated, aversive detail: <i>Tell me more about how this happened to you.</i> <i>Did it occur in the context of your job?</i></p>

Criterion B: Intrusion

Definition	Clarifying Prompts
<p>1. (B1) Repeated, disturbing, and unwanted memories of the stressful experience</p> <ul style="list-style-type: none"> • untriggered • while awake/not dreams • not rumination • not willfully thinking about it 	<p><i>How does it happen that you start remembering [EVENT]</i> <i>Do these memories occur while you are awake or asleep?</i> <i>Are these memories unwanted, or are you thinking of it on purpose?</i> <i>How much do these memories bother you?</i> <i>Are you able to put them out of your mind and think of something else?</i></p>
<p>2. (B2) Repeated, disturbing dreams of the stressful experience</p> <ul style="list-style-type: none"> • Dreams or nightmares • Content and/or affect related to the trauma • In children only, may be frightening dreams with unrecognizable content 	<p><i>Describe a typical dream. What happens?</i> <i>Do they wake you up?</i> <i>What do you experience when you wake up?</i> <i>How long does it take you to get back to sleep?</i> <i>How much sleep do you lose?</i> <i>How much do these dreams bother you?</i></p>

Definition	Clarifying Prompts
<p>3. (B3) Suddenly feeling or acting as if the stressful experience were actually happening again</p> <ul style="list-style-type: none"> • Differentiated from dreaming • Quality of a flashback or of reliving the experience • Some loss of connection to the present reality • May be momentary or prolonged • May be subtle or dramatic • If very severe may include images, sound, smells, physical responses 	<p><i>How much does it seem as if [EVENT] were happening again?</i></p> <p><i>Are you confused about where you actually are?</i></p> <p><i>What do you do while this is happening?</i></p> <p><i>Do other people notice your behavior?</i></p> <p><i>What do they say or do?</i></p> <p><i>How long does it last?</i></p>
<p>4. (B4) Feeling very upset when something reminded you of the stressful experience</p> <ul style="list-style-type: none"> • Distressing emotional reactions e.g., sad, angry, afraid, upset, ashamed • May be triggered by internal or external cues 	<p><i>What kinds of reminders make you upset?</i></p> <p><i>How do you feel when this happens?</i></p> <p><i>How much do these reminders bother you?</i></p> <p><i>Are you able to calm yourself down when this happens? How long does it take?</i></p>
<p>5. (B5) Having strong physical reactions when something reminded you of the stressful experience</p> <ul style="list-style-type: none"> • Distressing physiological reactions e.g. sweating, trembling, shaking, heart racing panting, trouble breathing • May be triggered by internal or external cues 	<p><i>Can you give me some examples?</i></p> <p><i>Does your heart race or your breathing change?</i></p> <p><i>What about sweating or feeling really tense or shaky?</i></p> <p><i>What kinds of reminders trigger these reactions?</i></p> <p><i>How long does it take you to recover?</i></p>

Criterion C. Avoidance

Definition	Clarifying Prompts
<p>6. (C1) Avoiding memories, thoughts, or feelings related to the trauma</p>	<p><i>What kinds of thoughts or feelings do you avoid?</i></p> <p><i>How hard do you try to avoid these thoughts or feelings?</i></p> <p><i>What kinds of things do you do?</i></p>
<p>7. (C2) Avoiding external reminders of the stressful trauma (e.g., people, places, conversations, activities, objects, or situations)</p>	<p><i>What kinds of things do you avoid?</i></p> <p><i>How much effort do you make to avoid these reminders?</i></p> <p><i>Do you have to plan or change your activities to avoid them?</i></p> <p><i>Overall, how much of a problem is this for you?</i></p>

Criterion D. Negative alteration of cognition and mood

Definition	Clarifying Prompts
<p>8. (D1) Trouble remembering important parts of the stressful experience</p> <ul style="list-style-type: none"> • Not due to drugs • Not due to alcohol • Not due to loss of consciousness • Not due to normal inattention or forgetting 	<p><i>Do you feel there are gaps in your memory of [EVENT]?</i> <i>What parts have you had difficulty remembering?</i> <i>Do you feel you should be able to remember these things?</i> <i>Why do you think you can't?</i> <i>Did you have a head injury during [EVENT]?</i> <i>Were you knocked unconscious? Were you intoxicated from alcohol or drugs?</i> <i>Is this just normal forgetting? Or do you think you may have blocked it out because it would be too painful to remember?</i></p>
<p>9. (D2) Having strong negative beliefs about yourself, other people, or the world (for example, having thoughts such as: I am bad, there is something seriously wrong with me, no one can be trusted, the world is completely dangerous)</p> <ul style="list-style-type: none"> • Beliefs or expectations • Exaggerated or persistent • Regarding the self, world or others • Began or got worse after the event 	<p><i>Can you give me some examples?</i> <i>What about believing things like "I am bad," "there is something seriously wrong with me," "no one can be trusted," "the world is completely dangerous"?</i> <i>How strong are these beliefs?</i> <i>How convinced are you that these beliefs are actually true?</i> <i>Can you see other ways of thinking about it?</i> <i>How much of the time in the past month have you felt that way?</i> <i>Did these beliefs start or get worse after (EVENT)?</i> <i>Do you think they are related to the event? How so?</i></p>
<p>10. (D3) Blaming yourself or someone else for the stressful experience or what happened after it</p> <ul style="list-style-type: none"> • Exaggerated or distorted cognitions about causes or consequences of trauma • Not realistic or reasonable (e.g., blame the perpetrator, or person who deliberately caused the event or intended harm) 	<p>(Query self-blame and other blame separately if both are present.)</p> <p><i>In what sense do you see [YOURSELF, OTHERS] as having caused [EVENT]?</i> <i>Is it because of something [YOU OR OTHERS] did?</i> <i>Or something you think [YOU OR OTHERS] should have done but didn't?</i> <i>Is it because of something about [YOU OR OTHERS] in general?</i> <i>How much do you blame (YOURSELF OR OTHERS)?</i> <i>How convinced are you that [YOU OR OTHERS] are truly responsible for what happened?</i> <i>Do other people agree with you?</i> <i>Can you see other ways of thinking about it?</i></p>
<p>11. (D4) Having strong negative feelings such as fear, horror, anger, guilt, or shame</p> <ul style="list-style-type: none"> • Persistent • Began or got worse after the event 	<p><i>Can you give me some examples?</i> <i>What negative feelings do you experience?</i> <i>How strong are these negative feelings?</i> <i>How well are you able to manage them?</i> <i>How much of the time in the past month have you felt that way?</i> <i>Did these negative feelings start or get worse after (EVENT)?</i></p>

Definition	Clarifying Prompts
<p>12. (D5) Loss of interest in activities that you used to enjoy</p> <ul style="list-style-type: none"> • Not due to lack of opportunity, physical inability, or developmentally appropriate change in preferred activities 	<p><i>What kinds of things have you lost interest in or don't do as much as you used to</i></p> <p><i>Anything else?</i></p> <p><i>Why is that?</i></p> <p><i>Would you still enjoy [ACTIVITIES] once you got started?</i></p> <p><i>How many of your usual activities have you been less interested in?</i></p> <p><i>What kinds of things do you still enjoy doing?</i></p> <p><i>Did this loss of interest start or get worse after (EVENT)?</i></p>
<p>13. (D6) Feeling distant or cut off from other people</p> <ul style="list-style-type: none"> • Feeling distant or cut off from others 	<p><i>Tell me more about that.</i></p> <p><i>How strong are your feelings of being distant or cut off from others?</i></p> <p><i>Who do you feel closest to?</i></p> <p><i>How many people do you feel comfortable talking with about personal things?</i></p> <p><i>How much of the time in the past month have you felt that way?</i></p> <p><i>Did this feeling of being distant or cut off start or get worse after (EVENT)?</i></p>
<p>14. (D7) Trouble experiencing positive feelings (for example, being unable to feel happiness or have loving feelings for people close to you)</p>	<p><i>Tell me more about that.</i></p> <p><i>What feelings are difficult to experience?</i></p> <p><i>How much difficulty do you have experiencing positive feelings?</i></p> <p><i>Are you still able to experience any positive feelings?</i></p> <p><i>How much of the time do you feel that way?</i></p> <p><i>Did this trouble experiencing positive feelings start or get worse after (EVENT)?</i></p>

Criterion E. Alterations in arousal and reactivity

Definition	Clarifying Prompts
<p>15. (E1) Irritable behavior, angry outbursts, or acting aggressively</p> <ul style="list-style-type: none"> • Behaviors • Irritable, angry feelings are not coded unless acted upon • Began or worsened after the trauma 	<p><i>Can you give me some examples?</i></p> <p><i>How do you show it?</i></p> <p><i>Do you raise your voice or yell?</i></p> <p><i>Throw or hit things?</i></p> <p><i>Push or hit other people?</i></p>
<p>16. (E2) Taking too many risks or doing things that could cause you harm</p>	<p><i>Can you give me some examples?</i></p> <p><i>How much of a risk do you take?</i></p> <p><i>How dangerous are these behaviors?</i></p> <p><i>Were you injured or harmed in some way?</i></p> <p><i>How often?</i></p> <p><i>Did this behavior start or get worse after (EVENT)?</i></p>

Definition	Clarifying Prompts
<p>17. (E3) Being “superalert” or watchful or on guard</p> <ul style="list-style-type: none"> Alert or watchful behavior in the absence of a specific threat or danger 	<p>Have you felt as if you had to be on guard? Can you give me some examples? What kinds of things do you do when you’re alert or watchful? What causes you to react this way? Do you feel like you’re in danger or threatened in some way? Do you feel that way more than most people would in the same situation? Did being especially alert or watchful start or get worse after [EVENT]?</p>
<p>18. (E4) Feeling jumpy or easily startled</p> <ul style="list-style-type: none"> Startle response is extreme compared to most people Recovery after startle is delayed compared to most people Began or worsened after the event 	<p>What kinds of things made you startle? How strong are these startle reactions? How strong are they compared to how most people would respond? Do you do anything other people would notice? How long does it take you to recover? Did these startle reactions start or get worse after (EVENT)?</p>
<p>19. (E5) Having difficulty concentrating</p> <ul style="list-style-type: none"> Began or worsened after the event <p>Moderate - some of the time (20-30%)/problem concentrating clearly present, some difficulty but can concentrate with effort</p> <p>Severe - much of the time (50-60%)/pronounced problem concentrating, considerable difficulty even with effort</p>	<p>Can you give me some examples? Are you able to concentrate if you really try? Did these problems with concentration start or get worse after (EVENT)?</p>
<p>20. (E6) Trouble falling or staying asleep</p> <ul style="list-style-type: none"> Began or worsened after the trauma <p>Moderate - 30-90-min. loss of sleep</p> <p>Severe - 90 min. - 3 hrs. loss of sleep</p> <p>Extreme - >3 hours loss of sleep</p>	<p>What kinds of problems? How long does it take you to fall asleep? How often do you wake up in the night? Do you wake up earlier than you want to? How many total hours do you sleep each night? How many hours do you think you should be sleeping? How often in the past month have you had these sleep problems? Did these sleep problems start or get worse after (EVENT)?</p>